

L I S T
OF THE
S P E C I M E N S
OF
HYMENOPTEROUS INSECTS
IN
THE COLLECTION
OF THE
BRITISH MUSEUM.

PART II. — CHALCIDITES.
ADDITIONAL SPECIES.

PRINTED BY ORDER OF THE TRUSTEES.
LONDON, 1848.

INTRODUCTION.

THE principal object of the present Catalogue has been to give a complete list of the specimens of Hymenopterous Insects contained in the collection of the British Museum, indicating at the same time the peculiarities of each, as regards variations in colour, locality, and the source from whence it has been derived.

With this view, the different individuals of each species contained in the collection are indicated by the letters *a*, *b*, *c*, &c., following the name of the species, and its synonymes.

In the adoption of generic and specific names, it has been thought right to use, wherever it was possible, that which was first applied to the species. As far as regards the specific names, there is comparatively little difficulty in the application of this simple rule; but generic names have been used by different authors in senses so widely different, and the groups which they were meant to designate have been so variously extended or restricted, that it is no easy matter to determine, where several names have been used, which of them ought to be preferred.

If the generic name adopted be different from that first employed for the species after the establishment of the Linnean system of nomenclature, the name under which it was first described follows as a synonyme, and where the Insect has received more than one specific name, those names are also given, as being sufficient for all purposes of identification.

These are followed by a reference to a few of the best original figures of the species.

In those cases where the two sexes of the same species, or any particular individual state or variety belonging to it, have been differently named, the name or figure exclusively appertaining to the state or individual is placed after the indication of the specimen to which it applies.

This part, containing the Chalciditous Insects of this order which have been procured chiefly from Mr. F. Walker since the former part was published, has been kindly prepared by that gentleman, who has devoted his attention to these Insects, and has been the original describer of the greater number of the species. The Appendix of this part, also written by Mr. Walker, contains the descriptions of the hitherto undescribed species contained in the collection.

The part containing the other families will follow as soon as they are prepared.

JOHN EDWARD GRAY.

British Museum,
April 20th, 1848.

LIST OF CHALCIDITES.

SUPPLEMENT.

Order.—CHALCIDITES.

Family. LEUCOSPIDÆ.

LEUCOSPIS, *Fabr.*

- LEUCOSPIS Cayennensis? *Westwood, Germar. Zeit. Ent. t. 4, f. 4.*
a. Isle St. Thomas. From Mr. Walker's collection.

Fam. CHALCIDÆ.

SMIERA, *Spinola.*

- SMIERA nigrifex, *Walker, Cat. Chalc. B. M. p. 2.*
l. Prussia. Presented by E. Doubleday, Esq.
m. Prussia. From Mr. Walker's collection.

CHALCIS, *Fabr.*

- CHALCIS intermedia, *Cat. Chalc. B. M. p. 4. Nees. Hym. Ich. aff.*
Mon. ii. 29, 8.
a. Prussia. Presented by E. Doubleday, Esq.
b. Prussia. From Mr. Walker's collection.

Fam. EURYTOMIDÆ.

EURYTOMA, *Ill.*

- EURYTOMA atra. *Isosoma atrum, Walker, Ent. Mag. i. 14.*
a. England. From Mr. Walker's collection.

EURYTOMA Cestius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

EURYTOMA Mucianus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

ISOSOMA, *Walker.*

ISOSOMA tenuicorne, *Walker, Ent. Mag.* i. 19.
a. England. From Mr. Walker's collection.

ISOSOMA pusillum, *Walker, Ent. Mag.* i. 20.
a. England. From Mr. Walker's collection.

ISOSOMA breviventre, *Walker, Ent. Mag.* i. 20.
a. England. From Mr. Walker's collection.

ISOSOMA vacillans, *Walker, Ent. Mag.* iv. 23.
a. England. From Mr. Walker's collection.

ISOSOMA Nepe, *Walker, Ann. Nat. Hist.* xiv. 21.
a. England. From Mr. Walker's collection.

DECATOMA, *Spin.*

DECATOMA plana, *Walker, Ent. Mag.* i. 27.
a. England. From Mr. Walker's collection.

DECATOMA tenuicornis, *Walker, Ent. Mag.* i. 28.
a. England. From Mr. Walker's collection.

DECATOMA mesomelas, *Walker, Ent. Mag.* iv. 24.
a. England. From Mr. Walker's collection.

DECATOMA flavicornis, *Walker, Ent. Mag.* iv. 25.
a. England. From Mr. Walker's collection.

DECATOMA Nicææ, *Walker, Ann. Nat. Hist.* xiv. 22.
a. England. From Mr. Walker's collection.

Fam. TORYMIDÆ.

MEGASTIGMUS, *Spin.*

MEGASTIGMUS dorsalis, *Cat. Chalc. B. M.* p. 14 *Torymus stigma*,
Nees. Hym. Ich. aff. Mon. ii. 65, 15. *T. giganteus*, var.?
Kollar.

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

MONODONTOMERUS, *Westw.*

MONODONTOMERUS stigma. *Torymus ater*, *Nees. Hym. Ich. aff. Mon. ii. 69, 21.*

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

MONODONTOMERUS cupreus. *Torymus cupreus*, *Nees. Hym. Ich. aff. Mon. ii. 67, 18.* *Monodontomerus æreus*, *De Laporte, Walker, Ent. Mag. ii. 158.*

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

MONODONTOMERUS obsoletus. *Torymus obsoletus*, *Nees. Hym. Ich. aff. Mon. ii. 68, 19.* ? *Dipl. obsoleta*, *Fabr. Syst. Piezat. 150, 10. Spin. Ins. Lig. Fasc. iii. 159, 8. Fasc. iv. 213. Ann. Mus. vii. 148.* *Cynips obsoleta*, *Latr. Gen. Cr. et Ins. iv. 28.* *Ich. obsoletus*, *Fabr. Ent. Syst. Suppl. 218, 230?* *Coqueb. Ill. con. i. 21. Tab. 5, f. 2.* *Ich. rodophthalmus*, *Rossi, Mant. i. 125, 276.*

a. Prussia. From Mr. Walker's collection.

CALLIMOME, *Spin.*

CALLIMOME cynipedis, *Cat. Chalc. B. M. p. 15.*

c. Prussia. From Mr. Walker's collection.

CALLIMOME cingulatus, *Cat. Chalc. B. M. p. 15.*

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

CALLIMOME regalis, *Walker, Ent. Mag. i. 119.*

a. England. From Mr. Walker's collection.

CALLIMOME Bedeguaris, *Cat. Chalc. B. M. p. 16.*

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

CALLIMOME elegans, *Cat. Chalc. B. M. p. 16.* *Torymus auronitens*, *Foerster, Mon. Pterom. i. xxx. 14.*

d. Prussia. From Mr. Walker's collection.

CALLIMOME nigricornis, *Cat. Chalc. B. M. p. 17.*

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

CALLIMOME mutabilis, *Cat. Chalc. B. M. p. 17.* *Torymus compressus*, *Foerster, Mon. Pterom. i. xxxii. 6.*

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

CALLIMOME regius, *Cat. Chalc. B. M.* p. 19. C. inconstans, *De Lap., &c.* Torymus regius, *Nees. Hym. Ich. aff. Mon.* ii. 55, 1. *De Geer, Ins.* ii. T. incertus, *Foerster, Mon. Pterom.* l. xxxi. 25, 187. *t.* 31, *f.* 21. T. pubescens, *Foerster, Mon. Pterom.* l. xxx. 15.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

CALLIMOME Rasaces, *Walker, Ann. Nat. Hist.* xiv. 181.

a. England. From Mr. Walker's collection.

CALLIMOME formosus, *Walker, Ent. Mag.* i. 122.

a. England. From Mr. Walker's collection.

CALLIMOME macropterus, *Walker, Ent. Mag.* i. 124.

a. England. From Mr. Walker's collection.

CALLIMOME nitens, *Walker, Ent. Mag.* i. 126.

a. England. From Mr. Walker's collection.

CALLIMOME brevicauda, *Walker, Ent. Mag.* i. 126.

a. England. From Mr. Walker's collection.

CALLIMOME microcerus, *Walker, Ent. Mag.* i. 128.

a. France. From Mr. Walker's collection.

CALLIMOME micropterus, *Walker, Ent. Mag.* i. 130.

a. England. From Mr. Walker's collection.

CALLIMOME viridiæneus, *Walker, Ent. Mag.* i. 131

a. England. From Mr. Walker's collection.

CALLIMOME moestus, *Walker, Ent. Mag.* i. 133.

a. England. From Mr. Walker's collection.

CALLIMOME stramineitarsus, *Walker, Ent. Mag.* i. 135.

a. England. From Mr. Walker's collection.

CALLIMOME ater, *Walker, Ent. Mag.* i. 138.

a. England. From Mr. Walker's collection.

CALLIMOME fuscipennis, *Walker, Ent. Mag.* i. 137.

a. England. From Mr. Walker's collection.

CALLIMOME nigritarsus, *Walker, Ent. Mag.* i. 134.

a. England. From Mr. Walker's collection.

CALLIMOME Angelicæ, *Walker, Ent. Mag.* iv. 25.

a. England. From Mr. Walker's collection.

CALLIMOME Ærope, *Walker, Ann. Nat. Hist.* xiv. 182.

a. England. From Mr. Walker's collection.

CALLIMOME meridionalis, *Walker, Ent. Mag.* i. 131.

a. France. From Mr. Walker's collection.

CALLIMOME arvernica, *Walker, Ent. Mag.* i. 134.

a. France. From Mr. Walker's collection.

CALLIMOME Euphorbiae, *De Laporte, Ent. Mag.* i. 134.

a. France. From Mr. Walker's collection.

CALLIMOME eurynotus. *Torymus eurynotus, Foerster, MSS.*

Walker, Ann. Nat. Hist.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

CALLIMOME chrysis. *Torymus chrysis, Nees. Hym. Ich. aff. Mon.*

ii. 67, 17.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

CALLIMOME Nepthys, *Walker, Ann. Nat. Hist.* 1848.

a. England. From Mr. Walker's collection.

CALLIMOME Solidus, *Walker.* See Appendix.

a. England. From Mr. Walker's collection.

ORMYRUS, *Westw.*

ORMYRUS punctiger, *Cat. Chalc. B. M.* p. 20. *Siphoneura vario-*

losa, Nees. Hym. Ich. aff. Mon. ii. 82, 1.

d. Prussia. Presented by E. Doubleday, Esq.

e. Prussia. From Mr. Walker's collection.

ORMYRUS cæruleus. *Siphonura cærulea, Foerster, MSS. Walker,*

Ann. Nat. Hist. 1848.

a. Prussia. From Mr. Walker's collection.

ORMYRUS nigrocyanus, *Walker. Ent. Mag.* i. 140.

a. England. From Mr. Walker's collection.

Fam. EUCHARIDÆ.

EUCHARIS, *Latr.*

EUCHARIS adscendens, *Cat. Chalc. B. M.* p. 21.

c. Germany. From Mr. Walker's collection.

PERILAMPUS, *Latr.*

PERILAMPUS micans, *Dalm. Kongl. Vet. Acad. Handl.* 1822.

a. Germany. From Mr. Walker's collection.

Fam. ————?

ELATUS, *Walker, MSS.* See Appendix.ELATUS Thenæ, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

Fam. SPALANGIIDÆ.

SPALANGIA, *Latr.*SPALANGIA nigra, *Cut. Chalc. B. M. p. 23.* S. nigripes. S. nigroænea, *Curtis, Brit. Ent. 140.*

e. England. From Mr. Walker's collection.

SPALANGIA hirta, *Haliday, Ent. Mag. i. 334.*

a. England. From Mr. Walker's collection.

CEROCEPHALA, *Westw.*CEROCEPHALA cornigera. C. cornigera, *Walker, Ent. Mag. ii. 149.*

a. Germany. From Mr. Walker's collection.

Fam. PTEROMALIDÆ.

ASAPHES, *Walker,*ASAPHES ænea. Chrysolampus suspensus, *Nees. Hym. Ich. aff. Mon. ii. 127.* C. altiventris, *Nees. Hym. &c. ii. 127.* Podagrion splendens? *Spin. Ann. Mus. vii. 147, 2.* Pteromalus petiolatus? *Zett. Ins. Lapp. xiv. 332.* Colax Aphidii, *Curtis, Journ. Roy. Agric. Soc. iii.*

a. England. From Mr. Walker's collection.

TETRAMESA, *Walker, MSS.* See Appendix.TETRAMESA Larbas, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

GASTRANCISTRUS, *Westw.*GASTRANCISTRUS viridis, *Walker, Ent. Mag. ii. 173.*

a. England. From Mr. Walker's collection.

GASTRANCISTRUS tenebriosus, *Walker, Ent. Mag. ii. 174.*

a. England. From Mr. Walker's collection.

GASTRANCISTRUS obscurellus, *Walker, Ent. Mag. ii. 17*

a. England. From Mr. Walker's collection.

- GASTRANCISTRUS Acontes, *Walker, Ann. Nat. Hist.* iv. 32.
a. Scotland. From Mr. Walker's collection.
- GASTRANCISTRUS longicornis, *Haliday, MSS.* See Appendix.
a. Ireland. From Mr. Walker's collection.
- GASTRANCISTRUS Dryas, *Walker, Mon. Chal.* i. 209.
a. South of France. From Mr. Walker's collection.
- GASTRANCISTRUS Panares, *Walker, Ann. Nat. Hist.* xiv. 339.
a. Alten, Finmark. From Mr. Walker's collection.
- GASTRANCISTRUS Amabœus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Accia, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Hamillus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Geganus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Fidenas, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Aleetus, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Hierocles, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Iriarte, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Amnisos, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Pacilus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- GASTRANCISTRUS Lœlianus, *Cat. Chalc. B. M.* See Appendix.
a. England. From Mr. Walker's collection.

PSILONOTUS, *Walker.*

- PSILONOTUS Adamas, *Walker, Ent. Mag.* ii. 179. *The Entomologist, pl. B. fig. 2.*
a. England. From Mr. Walker's collection.
- PSILONOTUS Hortensia, *Haliday, Ann. Nat. Hist.* 1846.
a. Ireland. From Mr. Walker's collection.
- PSILONOTUS Achæus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

ACRIAS, *Walker*.

ACRIAS Nileus, *Walker, Ann. Nat. Hist.* 1848, 29.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

RHAPHITELUS, *Walker*.

RHAPHITELUS maculatus, *Walker, Ent. Mag.* ii. 179. *The Entomologist*, pl. A, fig. 2. Pteromalus subulifer, *Foerster, Mon. Pterom.* i. xxviii. 264. Stylocerus subulifer, *Erichson, Bericht.* 1847.

a. England. From Mr. Walker's collection.

MACROMESUS, *Walker, MSS.* See Appendix.

MACROMESUS Amphiretus, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

MEROMALUS, *Walker*.

MEROMALUS flavicornis, *Walker, Ent. Mag.* ii. 178. Ormocerus Drymo, *Mon. Chal.* i. 205.

a. England. From Mr. Walker's collection.

MACROGLENES, *Westw.*

MACROGLENES penetrans, *Cat. Chal. B. M. p.* 24. *Trans. Ent. Soc.* iii. 295. *Curtis, Journ. Roy. Agric. Soc.* v.

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

PIRENE, *Haliday*.

PIRENE Scylax, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

PIRENE varicornis, *Hal. Trans. Ent. Soc.* iii. 296.

a. England. From Mr. Walker's collection.

PIRENE chalybea, *Hal. Trans. Ent. Soc.* iii. 296.

a. England. From Mr. Walker's collection.

PIRENE eximia, *Hal. Trans. Ent. Soc.* iii. 296.

a. England. From Mr. Walker's collection.

ORMOCERUS, *Walker*.

ORMOCERUS Sabbas, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

ORMOCERUS Aletes, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

ORMOCERUS Borges, *Walker, Mon. Chal.* i. 208.
a. Ireland. From Mr. Walker's collection.

ORMOCERUS Bambyce, *Walker, Mon. Chal.* i. 208.
a. England. From Mr. Walker's collection.

ORMOCERUS Vibulenus, *Walker, Mon. Chal.* i. 205.
a. England. From Mr. Walker's collection.

ORMOCERUS Mandrocles, *Walker, Mon. Chal.* i. 206.
a. England. From Mr. Walker's collection.

ORMOCERUS Pisuthrus, *Walker, Mon. Chal.* i. 207.
a. England. From Mr. Walker's collection.

UROLEPIS, *Walker.*

UROLEPIS maritima, *Cat. Chalc. B. M. p.* 26. *Pteromalus salinus?*
Von Heyden. Ent. Zeit. 1847.
b. England. From Mr. Walker's collection.

CYRTOGASTER, *Walker.*

CYRTOGASTER vulgaris, *Cat. Chalc. B. M. p.* 26. *Dicormus aquis-*
granensis, Foerster, Mon. Pterom. i. xxxvi.
c. Prussia. Presented by E. Doubleday, Esq.
d. Prussia. From Mr. Walker's collection.

CYRTOGASTER Scotica, *Walker, Ent. Mag.* i. 382.
a. Scotland. From Mr. Walker's collection.

CYRTOGASTER thoracica, *Walker, Ent. Mag.* i. 383.
a. England. From Mr. Walker's collection.

CYRTOGASTER pusilla, *Walker, Ent. Mag.* i. 383.
a. England. From Mr. Walker's collection.

CYRTOGASTER clavicornis, *Walker, Ent. Mag.* i. 383.
a. England. From Mr. Walker's collection.

CYRTOGASTER obscura, *Walker, Ent. Mag.* i. 383.
a. England. From Mr. Walker's collection.

CYRTOGASTER tenuis, *Walker, Ent. Mag.* i. 383.
a. England. From Mr. Walker's collection.

CYRTOGASTER cingulipes, *Walker, Ent. Mag.* i. 384.
a. England. From Mr. Walker's collection.

CYRTOGASTER Pæsos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

TITYROS, *Walker, MSS.* See Appendix.

TITYROS Poreia, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

PACHYLARTHURUS, *Westw.*

PACHYLARTHURUS flavicornis. Phacostomus patellanus, *Nees, Hym. Ich. aff. Mon. ii. 121.*

a. From Mr. Walker's collection.

PACHYLARTHURUS patellanus. Phacostomus similis, *Foerster, Mon. Pterom. i. xxxii. 2.* Pteromalus palpigerus, *Zetterstedt, Ins. Lapp. 3.*

a. Prussia. From Mr. Walker's collection.

PACHYLARTHURUS Elyces, *Walker, Mon. Chal. i. 195.*

a. France. From Mr. Walker's collection.

DICYCLUS, *Walker.*

DICYCLUS brevicornis, *Walker, Ent. Mag. i. 456.*

a. England. From Mr. Walker's collection.

TOXEUMA, *Walker.*

TOXEUMA Ericæ, *Cat. Chalc. B. M. p. 28.*

b. Prussia. Presented by E. Doubleday, Esq.

SPHEGIGASTER, *Spinola.*

SPHEGIGASTER flavicornis, *Cat. Chalc. B. M. p. 28.* Chrysolampus coronatus, *Foerster Mon. Pterom. i. 35, 5.*

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

SPHEGIGASTER clavicornis. Merismus clavicornis, *Walker, Ent. Mag. i. 377.*

a. England. From Mr. Walker's collection.

SPHEGIGASTER rufipes. Merismus rufipes, *Walker, Ent. Mag. i. 378.*

a. England. From Mr. Walker's collection.

SPHEGIGASTER Agriope, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

SPHEGIGASTER Lasthenes, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

SYNTOMOPUS, *Walker*.

SYNTOMOPUS thoracicus, *Walker, Ent. Mag.* i. 372.

a. England. From Mr. Walker's collection.

SYNTOMOPUS Dirce, *Walker. Miscogaster Dirce, Mon. Chal.*
i. 195.

a. England. From Mr. Walker's collection.

MEGORISMUS, *Walker*.

MEGORISMUS Aon, *Walker, Ann. Nat. Hist.* 1848.

a. From Mr. Walker's collection.

MEGORISMUS Nicææ *Miscogaster Nicææ, Walker, Mon. Chal.*
i. 197.

a. England. From Mr. Walker's collection.

CORYNA, *Walker*.

CORYNA clavata, *Cat. Chalc. B. M.* p. 29.

b. Prussia. Presented by E. Doubleday, Esq.

DIPARA, *Walker*.

DIPARA petiolata, *Walker, Ent. Mag.* i. 373.

a. England. From Mr. Walker's collection.

MERISUS, *Walker*.

MERISUS splendidus, *Walker, Ent. Mag.* ii. 167.

a. France. From Mr. Walker's collection.

MICROMELUS, *Walker*.

MICROMELUS pyrrhogaster. *Pteromalus mutilus, Foerster, Mon.*
Pterom. i. xxvii. 254.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

MERAPORUS, *Walker*.

MERAPORUS graminicola, *Cat. Chalc. B. M.* p. 35. *Curtis, Journ.*
Roy. Agric. Soc. vii. 97. *pl. P, fig.* 19.

b. England. From Mr. Walker's collection.

LAMPROTATUS, *Walker*.

LAMPROTATUS maculipes. *Miscogaster maculipes, Walker, Ent.*
Mag. i. 460. *Mon. Chal.* i. 282.

a. England. From Mr. Walker's collection.

- LAMPROTATUS æneus. *Miscogaster æneus*, *Walker, Ent. Mag. i.*
461. *Mon. Chal. i.* 285.
a. England. From Mr. Walker's collection.
- LAMPROTATUS nitidipes. *Miscogaster nitidipes*, *Walker, Ent.*
Mag. i. 462. *Mon. Chal. i.* 286.
a. England. From Mr. Walker's collection.
- LAMPROTATUS ovatus. *Miscogaster ovata*, *Walker, Ent. Mag. i.*
462. *Mon. Chal. i.* 287.
a. England. From Mr. Walker's collection.
- LAMPROTATUS convexus. *Miscogaster convexus*, *Walker, Ent. Mag.*
i. 463. *Mon. Chal. i.* 288.
a. England. From Mr. Walker's collection.
- LAMPROTATUS apicalis. *Miscogaster apicalis*, *Walker, Ent. Mag.*
i. 463. *Mon. Chal. i.* 288.
a. England. From Mr. Walker's collection.
- LAMPROTATUS tristis. *Miscogaster tristis*, *Walker, Ent. Mag. i.*
463. *Mon. Chal. i.* 288.
a. England. From Mr. Walker's collection.
- LAMPROTATUS costalis. *Miscogaster costalis*, *Walker, Ent. Mag.*
i. 463. *Mon. Chal. i.* 290.
a. England. From Mr. Walker's collection.
- LAMPROTATUS cyaneus. *Miscogaster cyanea*, *Walker, Ent. Mag. i.*
463. *Mon. Chal. i.* 290.
a. England. From Mr. Walker's collection.
- LAMPROTATUS contiguus. *Miscogaster contigua*, *Walker, Ent.*
Mag. i. 464. *Mon. Chal. i.* 291.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Rubrius, *Walker, Ann. Nat. Hist.* 1846, 111.
a. Scotland, From Mr. Walker's collection.
- LAMPROTATUS Helenor, *Walker, Ann. Nat. Hist.* 1846, 111.
a. Scotland. From Mr. Walker's collection.
- LAMPROTATUS Venilia, *Walker, Ann. Nat. Hist.* 1846, 113.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Phlegias, *Walker, Ann. Nat. Hist.* xiv. 332.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Mazæus, *Walker, Ann. Nat. Hist.* xiv. 333.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Brises, *Walker, Ann. Nat. Hist.* xiv. 333.
a. Alten, Finmark. From Mr. Walker's collection.

- LAMPROTATUS Cleta, *Walker, Ann. Nat. Hist.* xiv. 334.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Scœa, *Walker, Ann. Nat. Hist.* xiv. 335.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Leucon, *Walker, Ann. Nat. Hist.* xiv. 335.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Lesches, *Walker, Ann. Nat. Hist.* xiv. 336.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Icelos, *Walker, Ann. Nat. Hist.* xiv. 337.
a. Alten, Finmark. From Mr. Walker's collection.
- LAMPROTATUS Mallius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Zipoetes, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Œbares, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Phylander, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Acarnas, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Acilius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Amulius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Labaris, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Bolgius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Methymna, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Pycnos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- LAMPROTATUS Coretas, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

SELADERMA, *Walker.*

- SELADERMA breve, *Walker, Ent. Mag.* ii. 296.
a. England. From Mr. Walker's collection.

SELADERMA Capreæ. Cynips Capreæ, *Linn. Syst. Nat.* ii. 919, 14.
Faun. Suec. 1531. *Vill. Ent. Linn.* iii. 74, 14. *Fabr. Syst.*
Ent. ii. *Ent. Syst.* ii. 102, 13. *Syst. Piez.* 146, 15. *Pterom-*
malus Salicis, *Nees. Hym. Ich. aff. Mon.* ii. 105, 20.

a. England. From Mr. Walker's collection.

b. Prussia. From Mr. Walker's collection.

SELADERMA Saurus, *Walker, Ann. Nat. Hist.* xiv. 338.

a. Alten, Finmark. From Mr. Walker's collection.

SEMIOTUS, *Walker.*

SEMIOTUS Tauriscus, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

METASTENUS, *Walker.*

METASTENUS concinnus, *Walker, Ent. Mag.* ii. 302.

a. England. From Mr. Walker's collection.

PACHYNEURON, *Walker.*

PACHYNEURON Pruni, *Foerster, MSS. Walker, Ann. Nat. Hist.*
 1848.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

PACHYNEURON concolor, *Pteromalus concolor?* *Foerster, Mon.*
Pterom. i. xxvi. 236.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

MERAPORUS, *Walker.*

MERAPORUS exiguus, *Walker, Ent. Mag.* ii. 301.

a. England. From Mr. Walker's collection.

EUNOTUS, *Walker.*

EUNOTUS cretaceus, *Walker, Ent. Mag.* ii. 298. *The Entomologist*,
pl. A, fig. 4.

a. England. From Mr. Walker's collection.

SPANIOPUS, *Walker.*

SPANIOPUS dissimilis, *Walker, Ent. Mag.* i. 466. *Mon. Chal.*
 i. 292.

a. England. From Mr. Walker's collection.

PSILOCERA, *Walker*.

- PSILOCERA obscura, *Walker, Ent. Mag.* i. 374. *Metopon atrum*, *Fem. Ent. Mag.* ii. 303. *The Entomologist*, pl. B, fig. 3.
Merismus obscurus? *Blanchard, Goureau, Ann. Soc. Ent. Fr.*
2me. Série, iv. 478, pl. 11, f. 3.
 a. England. From Mr. Walker's collection.

PTEROMALUS, *Swederus*.

- PTEROMALUS fasciiventris, *Cat. Chalc. B. M.* p. 36.
 b. Prussia. Presented by E. Doubleday, Esq.
 PTEROMALUS tibialis, *Cat. Chalc. B. M.* p. 36. *P. sodalis*, *Foerster*,
Mon. Pterom. i. xi. 24.
 b. Prussia. From Mr. Walker's collection.
 PTEROMALUS dilectus, *Cat. Chalc. B. M.* p. 37.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
 PTEROMALUS catillus, *Cat. Chalc. B. M.* p. 40. *P. multicarinatus*,
Foerster, Mon. Pterom. i. xv. 83.
 b. Prussia. From Mr. Walker's collection.
 PTEROMALUS bracteatus, *Cat. Chalc. B. M.* p. 40. *P. chalcolampus*,
Foerster, Mon. Pterom. i. xvi. 103.
 b. Prussia. From Mr. Walker's collection.
 PTEROMALUS herbidus, *Cat. Chalc. B. M.* p. 40. *P. dilectus*,
Foerster, Mon. Pterom. i. xxiv. 205.
 b. Prussia. From Mr. Walker's collection.
 PTEROMALUS affinis, *Cat. Chalc. B. M.* p. 41. *P. statutus*, *Foerster*,
Mon. Pterom. i. xviii. 126.
 b. Prussia. From Mr. Walker's collection.
 PTEROMALUS fumipennis, *Cat. Chalc. B. M.* p. 41. *P. nubeculosus*,
Foerster, Mon. Pterom. i. xxviii. 260.
 b. Prussia. From Mr. Walker's collection.
 PTEROMALUS microcerus, *Cat. Chalc. B. M.* p. 42. *P. reconditus*,
Foerster, Mon. Pterom. i. xi. 25.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
 PTEROMALUS subniger, *Cat. Chal. B. M.* p. 42. *P. subniger*,
Foerster, Mon. Pterom. i. xx. 150.
 b. Prussia. From Mr. Walker's collection.

PTEROMALUS muscarum, *Cat. Chalc. p.* 42. *P. psittacinus*, *Foerster, Mon. Pterom. i. xv.* 86. *P. larvarum*, *Nees. Hym. Ich. aff. Mon. ii.* 95, 3. *P. Thessalus*, *Walker, Mon. Chal. i.* 268. *Diplolepis larvarum*, *Spin. Ins. Lig. Fasc. iii.* 162, 16.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

PTEROMALUS puparum, *Cat. Chalc. B. M. p.* 43.

d. Prussia. Presented by E. Doubleday, Esq.

e. Prussia. From Mr. Walker's collection.

PTEROMALUS mesochlorus, *Cat. Chalc. B. M. p.* 43. *P. acuminatus*, *Foerster, Mon. Pterom. i. xiv.* 70.

b. Prussia. From Mr. Walker's collection.

PTEROMALUS micans, *Cat. Chalc. B. M. p.* 43. *P. varians*, *Nees. Hym. Ich. aff. Mon. ii.* 101, 15.

c. Prussia. Presented by E. Doubleday, Esq.

PTEROMALUS divisus, *Cat. Chalc. B. M. p.* 44.

c. Prussia. From Mr. Walker's collection.

PTEROMALUS hilaris, *Cat. Chalc. B. M. p.* 45. *P. quæsitus*, *Foerster, Mon. Pterom. i. xi.* 30.

b. Prussia. From Mr. Walker's collection.

PTEROMALUS Mutia, *Cat. Chalc. B. M. p.* 46. *P. vorax*, *Foerster, Mon. Pterom. i. xiv.* 74.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

PTEROMALUS Lucilla, *Cat. Chalc. B. M. p.* 47. *P. foveolatus*, *Foerster, Mon. Pterom. i. xiii.* 48.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

PTEROMALUS terminalis. *Platyterma terminale*, *Walker, Ent. Mag. ii.* 306.

a. England. From Mr. Walker's collection.

PTEROMALUS remotus. *Platyterma remotum*, *Walker, Ent. Mag. ii.* 342.

a. England. From Mr. Walker's collection.

PTEROMALUS laticornis. *Platyterma laticorne*, *Walker, Ent. Mag. ii.* 304.

a. England. From Mr. Walker's collection.

PTEROMALUS fulvipennis. *Amblymerus fulvipennis*, *Walker, Ent. Mag. ii.* 344.

a. England. From Mr. Walker's collection.

- PTEROMALUS humilis.* *Amblymerus humilis*, *Walker, Ent. Mag.* ii. 346.
a. England. From Mr. Walker's collection.
- PTEROMALUS nitescens.* *Amblymerus nitescens*, *Walker, Ent. Mag.* ii. 347.
a. England. From Mr. Walker's collection.
- PTEROMALUS hebes.* *Amblymerus hebes*, *Walker, Ent. Mag.* ii. 348.
a. England. From Mr. Walker's collection.
- PTEROMALUS tenellus.* *Amblymerus tenellus*, *Walker, Ent. Mag.* ii. 348.
a. England. From Mr. Walker's collection.
- PTEROMALUS fulvipes.* *Amblymerus fulvipes*, *Walker, Ent. Mag.* ii. 348.
a. England. From Mr. Walker's collection.
- PTEROMALUS nanus.* *Amblymerus nanus*, *Walker, Ent. Mag.* ii. 349.
a. England. From Mr. Walker's collection.
- PTEROMALUS iners.* *Amblymerus iners*, *Walker, Ent. Mag.* ii. 50.
a. England. From Mr. Walker's collection.
- PTEROMALUS trossulus.* *Amblymerus trossulus*, *Walker, Ent. Mag.* ii. 350.
a. England. From Mr. Walker's collection.
- PTEROMALUS tenebricus.* *Amblymerus tenebricus*, *Walker, Ent. Mag.* ii. 351.
a. England. From Mr. Walker's collection.
- PTEROMALUS mirus.* *Amblymerus mirus*, *Walker, Ent. Mag.* ii. 350.
a. England. From Mr. Walker's collection.
- PTEROMALUS pygmæus.* *Eutelus pygmæus*, *Walker, Ent. Mag.*
a. England. From Mr. Walker's collection.
- PTEROMALUS ocellus.* *Eutelus ocellus*, *Walker, Ent. Mag.* ii. 359.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS catenatus.* *Eutelus catenatus*, *Walker, Ent. Mag.* ii. 362.
a. England. From Mr. Walker's collection.
- PTEROMALUS helvipes.* *Eutelus helvipes*, *Walker, Ent. Mag.* ii. 365.
a. England. From Mr. Walker's collection.

- PTEROMALUS politus*. *Eutelus politus*, *Walker, Ent. Mag.* ii. 369.
a. England. From Mr. Walker's collection.
- PTEROMALUS vagans*. *Eutelus vagans*, *Walker, Ent. Mag.* ii. 369.
a. England. From Mr. Walker's collection.
- PTEROMALUS perversus*, *Walker, Ent. Mag.* ii. 479.
a. England. From Mr. Walker's collection.
- PTEROMALUS extentus*, *Walker, Ent. Mag.* ii. 480.
a. England. From Mr. Walker's collection.
- PTEROMALUS megachlorus*, *Walker, Ent. Mag.* ii. 486.
a. England. From Mr. Walker's collection.
- PTEROMALUS aurifer*, *Walker, Ent. Mag.* ii. 487.
a. England. From Mr. Walker's collection.
- PTEROMALUS apertus*, *Walker, Ent. Mag.* ii. 489. *P. chalcophanes*,
Foerster, Mon. Pterom. i. xxv. 220.
a. England. From Mr. Walker's collection.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS cuprinus*, *Walker, Ent. Mag.* ii. 489.
a. England. From Mr. Walker's collection.
- PTEROMALUS curtus*, *Walker, Ent. Mag.* ii.
a. England. From Mr. Walker's collection.
- PTEROMALUS pinguis*, *Walker, Ent. Mag.* ii. 490.
a. England. From Mr. Walker's collection.
- PTEROMALUS despectus*, *Walker, Ent. Mag.* ii. 491.
a. England. From Mr. Walker's collection.
- PTEROMALUS purpureus*, *Walker, Ent. Mag.* ii. 493.
a. England. From Mr. Walker's collection.
- PTEROMALUS anticus*, *Walker, Ent. Mag.* ii. 494.
a. England. From Mr. Walker's collection.
- PTEROMALUS rufinus*, *Walker, Ent. Mag.* ii. 495.
a. England. From Mr. Walker's collection.
- PTEROMALUS saturatus*, *Walker, Ent. Mag.* ii. 495.
a. England. From Mr. Walker's collection.
- PTEROMALUS futilis*, *Walker, Ent. Mag.* ii. 496. *P. operosus*,
Foerster, Mon. Pterom. i. xi. 28.
a. England. From Mr. Walker's collection.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS ovatus*, *Nees, Hym. Ich. aff. Mon.* ii. 103, 18.
a. Prussia. From Mr. Walker's collection.

- PTEROMALUS prætermisus*, *Foerster, Mon. Pterom. i. xvii. 107.*
a. Prussia. From Mr. Walker's collection.
- PTEROMALUS inclytus*, *Foerster, Mon. Pterom. i. xix. 145.*
a. Prussia. From Mr. Walker's collection.
- PTEROMALUS tibialis*, *Foerster, MSS.*
a. Prussia. Presented by E. Doubleday, Esq.
- PTEROMALUS laticeps*, *Foerster, MSS.*
a. Prussia. Presented by E. Doubleday, Esq.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS distinguendus*, *Foerster, Mon. Pterom. i. xv. 84.*
a. Prussia. Presented by E. Doubleday, Esq.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS sulphuripes*, *Foerster, MSS.*
a. Prussia. From Mr. Walker's collection.
- PTEROMALUS sparsus*, *Foerster, Mon. Pterom. i. x. 18.*
a. Prussia. Presented by E. Doubleday, Esq.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS communis?* *Nees. Hym. Ich. aff. Mon. ii. 103, 17.*
a. Prussia. Presented by E. Doubleday, Esq.
- PTEROMALUS pexatus*, *Walker, Ent. Mag. ii. 499.*
a. England. From Mr. Walker's collection.
- PTEROMALUS detritus*, *Walker, Ent. Mag. ii. 499.*
a. England. From Mr. Walker's collection.
- PTEROMALUS latifrons*, *Walker, Ent. Mag. ii. 501.*
a. England. From Mr. Walker's collection.
- PTEROMALUS latipennis*, *Walker, Ent. Mag. iii. 96.*
a. England. From Mr. Walker's collection.
- PTEROMALUS mediocris*, *Walker, Ent. Mag. iii. 97.*
a. England. From Mr. Walker's collection.
- PTEROMALUS basalis*, *Walker, Ent. Mag. iii. 185.*
a. England. From Mr. Walker's collection.
- PTEROMALUS decisis*, *Walker, Ent. Mag. iii. 185.* *P. opulentus*,
Foerster, Mon. Pterom. i. xxiv. 213.
a. England. From Mr. Walker's collection.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS lautus*, *Walker, Ent. Mag. iii. 186.*
a. England. From Mr. Walker's collection.
- PTEROMALUS infectus*, *Walker, Ent. Mag. iii. 186.*
a. England. From Mr. Walker's collection.

- PTEROMALUS impeditus*, *Walker, Ent. Mag.* iii. 187.
a. England. From Mr. Walker's collection.
- PTEROMALUS grandiclava*, *Walker, Ent. Mag.* iii. 193.
a. England. From Mr. Walker's collection.
- PTEROMALUS fucicola*, *Walker, Ent. Mag.* iii. 194.
a. England. From Mr. Walker's collection.
- PTEROMALUS plenus*, *Walker, Ent. Mag.* iii. 199.
a. England. From Mr. Walker's collection.
- PTEROMALUS solutus*, *Walker, Ent. Mag.* iii. 199.
a. England. From Mr. Walker's collection.
- PTEROMALUS berylli*, *Walker, Ent. Mag.* iii. 199. *P. crassus*,
Foerster. Mon. Pterom. i. xxv. 218.
a. England. From Mr. Walker's collection.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS thoracicus*, *Walker, Ent. Mag.* iii. 200.
a. England. From Mr. Walker's collection.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS omnivorus*, *Walker, Ent. Mag.* iii. 204.
a. England. From Mr. Walker's collection.
b. Prussia. From Mr. Walker's collection.
- PTEROMALUS nigroæneus*, *Walker, Ent. Mag.* iii. 206.
a. England. From Mr. Walker's collection.
- PTEROMALUS cliens*, *Walker, Ent. Mag.* iii. 469.
a. England. From Mr. Walker's collection.
- PTEROMALUS illudens*, *Walker, Ent. Mag.* iii. 470.
a. England. From Mr. Walker's collection.
- PTEROMALUS gracilis*, *Walker, Ent. Mag.* iii. 477.
a. England. From Mr. Walker's collection.
- PTEROMALUS dorsalis*, *Walker, Ent. Mag.* iii. 478.
a. England. From Mr. Walker's collection.
- PTEROMALUS subquadratus*, *Walker, Ent. Mag.* iii. 478.
a. England. From Mr. Walker's collection.
- PTEROMALUS signatus*, *Walker, Ent. Mag.* iii. 479.
a. England. From Mr. Walker's collection.
- PTEROMALUS servulus*, *Walker, Ent. Mag.* iii. 481.
a. England. From Mr. Walker's collection.
- PTEROMALUS rusticus*, *Walker, Ent. Mag.* iii. 482.
a. England. From Mr. Walker's collection.

- PTEROMALUS diversus*, *Walker, Ent. Mag.* iii. 483.
a. England. From Mr. Walker's collection.
- PTEROMALUS thalassinus*, *Walker, Ent. Mag.* iii. 485.
a. England. From Mr. Walker's collection.
- PTEROMALUS confinis*, *Walker, Ent. Mag.* iii. 486.
a. England. From Mr. Walker's collection.
- PTEROMALUS exilis*, *Walker, Ent. Mag.* iii. 487.
a. England. From Mr. Walker's collection.
- PTEROMALUS chrysammos*, *Walker, Ent. Mag.* iii. 487.
a. England. From Mr. Walker's collection.
- PTEROMALUS pervasus*, *Walker, Ent. Mag.* iii. 489.
a. England. From Mr. Walker's collection.
- PTEROMALUS longulus*, *Walker, Ent. Mag.* iii. 491.
a. England. From Mr. Walker's collection.
- PTEROMALUS comes*, *Walker, Ent. Mag.* iii. 492.
a. England. From Mr. Walker's collection.
- PTEROMALUS gentilis*, *Walker, Ent. Mag.* iii. 493.
a. England. From Mr. Walker's collection.
- PTEROMALUS equestris*, *Walker, Ent. Mag.* iii. 495.
a. England. From Mr. Walker's collection.
- PTEROMALUS simulans*, *Walker, Ent. Mag.* iii. 495.
a. England. From Mr. Walker's collection.
- PTEROMALUS Empoclus*, *Walker, Mon. Chal.* i. 214.
a. England. From Mr. Walker's collection.
- PTEROMALUS Docimus*, *Walker, Mon. Chal.* i. 217.
a. England. From Mr. Walker's collection.
- PTEROMALUS Pronax*, *Walker, Mon. Chal.* i. 220.
a. England. From Mr. Walker's collection.
- PTEROMALUS Samus*, *Walker, Mon. Chal.* i. 221.
a. England. From Mr. Walker's collection.
- PTEROMALUS Vibullius*, *Walker, Mon. Chal.* i. 221.
a. England. From Mr. Walker's collection.
- PTEROMALUS Calligetis*, *Walker, Mon. Chal.* i. 222.
a. England. From Mr. Walker's collection.
- PTEROMALUS Japis*, *Walker, Mon. Chal.* i. 222.
a. England. From Mr. Walker's collection.
- PTEROMALUS Carcinus*, *Walker, Mon. Chal.* i. 229.
a. England. From Mr. Walker's collection.

- PTEROMALUS Gargasus, *Walker, Mon. Chal.* i. 229.
a. England. From Mr. Walker's collection.
- PTEROMALUS Flavius, *Walker, Mon. Chal.* i. 230.
a. England. From Mr. Walker's collection.
- PTEROMALUS Gonatas, *Walker, Mon. Chal.* i. 231.
a. England. From Mr. Walker's collection.
- PTEROMALUS Druso, *Walker, Mon. Chal.* i. 232.
a. England. From Mr. Walker's collection.
- PTEROMALUS Aper, *Walker, Mon. Chal.* i. 234.
a. England. From Mr. Walker's collection.
- PTEROMALUS Tigasis, *Walker, Mon. Chal.* i. 232.
a. England. From Mr. Walker's collection.
- PTEROMALUS Vala, *Walker, Mon. Chal.* i. 234.
a. England. From Mr. Walker's collection.
- PTEROMALUS Styrius, *Walker, Mon. Chal.* i. 243.
a. England. From Mr. Walker's collection.
- PTEROMALUS Ariovistus, *Walker, Mon. Chal.* i. 244.
a. England. From Mr. Walker's collection.
- PTEROMALUS Boreus, *Walker, Mon. Chal.* i. 246.
a. England. From Mr. Walker's collection.
- PTEROMALUS Hippos, *Walker, Mon. Chal.* i. 247.
a. England. From Mr. Walker's collection.
- PTEROMALUS Sophax, *Walker, Mon. Chal.* i. 249.
a. England. From Mr. Walker's collection.
- PTEROMALUS Faustina, *Walker, Mon. Chal.* i. 251.
a. England. From Mr. Walker's collection.
- PTEROMALUS Tiburtus, *Walker, Mon. Chal.* i. 251.
a. England. From Mr. Walker's collection.
- PTEROMALUS Abila, *Walker, Mon. Chal.* i. 252.
a. England. From Mr. Walker's collection.
- PTEROMALUS Automedon, *Walker, Mon. Chal.* i. 254.
a. England. From Mr. Walker's collection.
- PTEROMALUS Tereus, *Walker, Mon. Chal.* i. 255.
a. England. From Mr. Walker's collection.
- PTEROMALUS Nyctimus, *Walker, Mon. Chal.* i. 257.
a. England. From Mr. Walker's collection.
- PTEROMALUS Crotopus, *Walker, Mon. Chal.* i. 258.
a. England. From Mr. Walker's collection.

- PTEROMALUS Dercyllus, *Walker, Mon. Chal.* i. 259.
a. England. From Mr. Walker's collection.
- PTEROMALUS Archidemus, *Walker, Mon. Chal.* i. 259.
a. England. From Mr. Walker's collection.
- PTEROMALUS Orsippus, *Walker, Mon. Chal.* i. 260.
a. England. From Mr. Walker's collection.
- PTEROMALUS Aglaope, *Walker, Mon. Chal.* i. 261.
a. England. From Mr. Walker's collection.
- PTEROMALUS Erasippus, *Walker, Mon. Chal.* i. 261.
a. England. From Mr. Walker's collection.
- PTEROMALUS Fontanus, *Walker, Mon. Chal.* i. 262.
a. England. From Mr. Walker's collection.
- PTEROMALUS Favorinus, *Walker, Mon. Chal.* i. 263.
a. England. From Mr. Walker's collection.
- PTEROMALUS Coryphe, *Walker, Mon. Chal.* i. 266.
a. England. From Mr. Walker's collection.
- PTEROMALUS Acræa, *Walker, Mon. Chal.* i. 266.
a. England. From Mr. Walker's collection.
- PTEROMALUS Ormenus, *Walker, Mon. Chal.* i. 269.
a. England. From Mr. Walker's collection.
- PTEROMALUS Sophron, *Walker, Mon. Chal.* i. 270.
a. England. From Mr. Walker's collection.
- PTEROMALUS Zelus, *Walker, Mon. Chal.* i. 270.
a. England. From Mr. Walker's collection.
- PTEROMALUS Hecato, *Walker, Mon. Chal.* i. 271.
a. England. From Mr. Walker's collection.
- PTEROMALUS Phyllus, *Walker, Mon. Chal.* i. 272.
a. England. From Mr. Walker's collection.
- PTEROMALUS Cælius, *Walker, Mon. Chal.* i. 272.
a. England. From Mr. Walker's collection.
- PTEROMALUS Promulus, *Walker, Ann. Nat. Hist.* iv. 232.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS Varro, *Walker, Ann. Nat. Hist.* iv. 233.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS Rhinthon, *Walker, Ann. Nat. Hist.* xiv. 341.
a. Isle of Soröc, Finmark. From Mr. Walker's collection.
- PTEROMALUS Musæus, *Walker, Ann. Nat. Hist.* xiv. 340.
a. Alten, Finmark. From Mr. Walker's collection.

- PTEROMALUS Mazaces*, *Walker, Ann. Nat. Hist.* xiv. 341.
a. Alten, Finmark. From Mr. Walker's collection.
- PTEROMALUS Lausus*, *Walker, Ann. Nat. Hist.* xix. 392, 1.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Cratylus*, *Walker, Ann. Nat. Hist.* xix. 392, 2.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Cassotis*, *Walker, Ann. Nat. Hist.* xix. 393, 3.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Euthymus*, *Walker, Ann. Nat. Hist.* xix. 393, 4.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Dipsas*, *Walker, Ann. Nat. Hist.* xix. 394, 5.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Hermeas*, *Walker, Ann. Nat. Hist.* xix. 394, 6.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Epicles*, *Walker, Ann. Nat. Hist.* xix. 394, 7.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Damo*, *Walker, Ann. Nat. Hist.* xix. 395, 8.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Cæax*, *Walker, Ann. Nat. Hist.* xix. 395, 9.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Doryssus*, *Walker, Ann. Nat. Hist.* xix. 395, 10.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Timæa*, *Walker, Ann. Nat. Hist.* xix. 396, 11.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Orontas*, *Walker, Ann. Soc. Ent.* xix. 396, 12.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Dymnus*, *Walker, Ann. Nat. Hist.* xix. 397, 13.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Eryx*, *Walker, Ann. Nat. Hist.* xix. 397, 14.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Hybreas*, *Walker, Ann. Nat. Hist.* xix. 397, 15.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Eurypon*, *Walker, Ann. Nat. Hist.* xix. 398, 16.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- PTEROMALUS Acrotatus*, *Walker, Trans. Linn. Soc.* xx. 153.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS Bubaris*, *Walker, Trans. Linn. Soc.* xx. 154.
a. Scotland. From Mr. Walker's collection.

- PTEROMALUS Ection, *Walker, Trans. Linn. Soc.* xx. 154.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS Xanthe, *Walker, Trans. Linn. Soc.* xx. 155.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS Aollius, *Walker, Trans. Linn. Soc.* xx. 156.
a. North Wales. From Mr. Walker's collection.
- PTEROMALUS Antho, *Walker, Trans. Linn. Soc.* xx. 156.
a. England. From Mr. Walker's collection.
- PTEROMALUS Learchus, *Walker, Trans. Linn. Soc.* xx. 157.
a. Scotland. From Mr. Walker's collection.
- PTEROMALUS Autorides, *Walker, Trans. Linn. Soc.* xx. 157.
a. England. From Mr. Walker's collection.
- PTEROMALUS Saravus, *Walker, Trans. Linn. Soc.* xx. 157.
a. England. From Mr. Walker's collection.
- PTEROMALUS Anaxenor, *Walker, Trans. Linn. Soc.* xx. 158.
a. England. From Mr. Walker's collection.
- PTEROMALUS Tedanius, *Walker, Trans. Linn. Soc.* xx. 158.
a. England. From Mr. Walker's collection.
- PTEROMALUS Naubolus, *Walker, Trans. Linn. Soc.* xx. 159. P.
 larvarum? *Nees. Hym. Ich. aff. Mon.* ii. 94, 422. Dipl.
 lepis larvarum? *Spin. Ins. Lig. Fasc.* iii. 162, 16. Dipl.
 Braconidis? *Bouché, Natur. Gesch. der. Ins.* i. 171, 65.
a. England. From Mr. Walker's collection.
- PTEROMALUS Aglaus, *Walker, Trans. Linn. Soc.* xx. 160.
a. England. From Mr. Walker's collection.
- PTEROMALUS Urgo, *Walker, Trans. Linn. Soc.* xx. 160.
a. England. From Mr. Walker's collection.
- PTEROMALUS Orinus, *Walker, Trans. Linn. Soc.* xx. 161.
a. England. From Mr. Walker's collection.
- PTEROMALUS Æson, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Camma, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Phasis, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Cyamon. See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Hylœ, *Walker.* See Appendix.
a. England. From Mr. Walker's collection.

- PTEROMALUS Acco, *Walker*. See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Tychon, *Walker*. See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Cerycus, *Walker*. See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Arestor, *Walker*. See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Eulimene, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Dipænos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Glautias, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Suia, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Amnisos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Zagreus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Opheltes, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Mese, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Alopheus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Phylacis, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Cæno, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Corion, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Hermachus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Temesa, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Drepanon, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

- PTEROMALUS Belesis, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Lebadeia, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Berycynthos, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Lebenc, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Cercaphrus, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Cytæum, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Lampé, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Anchinoë, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Gallonius, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Eleuthera, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Lyttus, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Bienna, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Patro, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Lissos, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Morys, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Elyros, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Rhytium, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Myle, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.
- PTEROMALUS Loeta, *Walker, MSS.* See Appendix.
 a. England. From Mr. Walker's collection.

- PTEROMALUS** Androbius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Inatos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Axos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Otos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Carma, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Sybritia, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Mesapos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Larymna, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Etearchus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Phalasurna, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Themiso, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Allutius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Priansos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Gigon, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Orthagus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Diomedon, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Alimentus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Cœlius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS** Æacus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

- PTEROMALUS Alope**, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Cepio**, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Hyrtacina**, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Leuce**, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- PTEROMALUS Hyrtacus**, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

HETROXYS, *Westwood.*

- HETROXYS invenustus.** *Pteromalus invenustus, Walker, Ent. Mag.*
iv. 11.
a. England. From Mr. Walker's collection.
- HETROXYS macromerus.** *Pteromalus macromerus. Walker, Ent.*
Mag. iv. 11.
a. England. From Mr. Walker's collection.
- HETROXYS præpileus.** *Pteromalus præpileus, Walker, Ent. Mag.*
iv. 12.
a. England. From Mr. Walker's collection.
- HETROXYS dimidiatus.** *Pteromalus dimidiatus, Walker, Ent.*
Mag. iv. 12.
a. England. From Mr. Walker's collection.
- HETROXYS Aponius.** *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

CHEIROPACHUS, *Westwood.*

- CHEIROPACHUS Colon.** *Sphex Colon, Linn. Syst. Nat. i. 2, 945,*
28. Faun. Suec. 1656. Diplolepis quadrum, Fabr. Ptero-
malus quadrum, Zetterstedt, Ins. Lapp. 421. Cheiropachus
quadrum, Cat. Chalc. B. M. p. 50.
c. Prussia. Presented by E. Doubleday, Esq.
d. Prussia. From Mr. Walker's collection.

MACRONEURA, *Walker.*

- MACRONEURA maculipes**, *Cat. Chalc. B. M. p. 51.*
b. Prussia. Presented by E. Doubleday, Esq.

TRIGONODERUS, *Westwood.*

TRIGONODERUS tristis, *Walker, Ent. Mag.* iv. 17.

a. England. From Mr. Walker's collection.

TRIGONODERUS amabilis, *Walker, Ent. Mag.* iv. 20.

a. England. From Mr. Walker's collection.

TRIGONODERUS contemptus, *Walker, Ent. Mag.* iv. 22.

a. England. From Mr. Walker's collection.

TRIGONODERUS atrovirens, *Walker, Ent. Mag.* iv. 22.

a. England. From Mr. Walker's collection.

TRIGONODERUS Lappa, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

TRIGONODERUS Tarrha, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

TRIGONODERUS Alebion, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

TRIGONODERUS Polichna, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

PLATYNOCHEILUS, *Westwood.*

PLATYNOCHEILUS Erichsonii, *Westwood, Ent. Mag.* iii. 437. *Stenocera Erichsonii, Walker, Mon. Chal.* i. 276.

a. England. From Mr. Walker's collection.

PLATYNOCHEILUS Derceto. *Stenocera Derceto, Walker, Mon. Chal.* i. 275.

a. England. From Mr. Walker's collection.

STENOCERA, *Walker.*

STENOCERA Walkeri, *Curtis, Brit. Ent.* 596. *Walker, Ent. Mag.* iv. 357.

a. England. From Mr. Walker's collection.

NOTANISUS, *Walker.*

NOTANISUS versicolor, *Walker, Ent. Mag.* iv. 352.

a. South of France. From Mr. Walker's collection.

CLEONYMUS, *Latr.*

CLEONYMUS obscurus, *Walker, Ent. Mag.* iv. 352.

a. England. From Mr. Walker's collection.

EUELMUS, *Dalman*.

EUELMUS Degeeri, *Cat. Chalc. B. M.* p. 53.

d. Prussia. Presented by E. Doubleday, Esq.

EUELMUS Fonteia, *Walker, Ann. Nat. Hist.* xx. 19.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EUELMUS Lamachus, *Walker, Ann. Nat. Hist.* xx. 20.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EUELMUS, Epicaste, *Walker, Ann. Nat. Hist.* xx. 20.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EUELMUS Acron, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

EUELMUS excavatus, *Dalm. Kongl. Vet. Acad. Handl.* 1820.

Nees, ab. Ess. Hym. Ich. aff. Mon. ii. 79. *Walker, Ent. Mag.*

iv. 362. *Urocryptus excavatus, Westwood.*

a. England. From Mr. Walker's collection.

ERICYDNUS, *Haliday*.

ERICYDNUS strigosus, *Cat. Chalc. B. M.* p. 53.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

CHEILONEURUS, *Westwood*.

CHEILONEURUS elegans, *Cat. Chalc. B. M.* p. 54.

a. Prussia. Presented by E. Doubleday, Esq.

METALLON, *Walker, MSS.*

METALLON Acacallis, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

CERCHYSIUS, *Westwood*.

CERCHYSIUS Vulso, *Walker, Ann. Nat. Hist.* xix. 114.

a. England. From Mr. Walker's collection.

CERCHYSIUS Flaccus, *Walker, Ann. Nat. Hist.* xx. 21.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

ENCYRTUS, *Dalman*.

ENCYRTUS cyaneus, *Cat. Chalc. B. M.* p. 54.

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

- ENCYRTUS *Glaphyra*, *Cat. Chalc. B. M.* p. 55.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS *barbarus*, *Cat. Chalc. B. M.* p. 55.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS *subplanus*, *Cat. Chalc. B. M.* p. 56. E. *sceptriger*,
Foerster, Mon. Pterom. i. xlv. 3.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS *truncatellus*, *Cat. Chalc. B. M.* p. 56.
 d. Prussia. From Mr. Walker's collection.
- ENCYRTUS *Swederi*, *Cat. Chalc. B. M.* p. 56.
 d. Prussia. Presented by E. Doubleday, Esq.
 e. Prussia. From Mr. Walker's collection.
- ENCYRTUS *tessellatus*, *Cat. Chalc. B. M.* p. 57.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS *sericeus*, *Cat. Chalc. B. M.* p. 57.
 d. Prussia. Presented by E. Doubleday, Esq.
 e. Prussia. From Mr. Walker's collection.
- ENCYRTUS *punctipes*, *Cat. Chalc. B. M.* p. 57.
 c. Prussia. From Mr. Walker's collection.
 d. Prussia. From Mr. Walker's collection.
- ENCYRTUS *fuscicollis*, *Cat. Chalc. B. M.* p. 58.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. England. From Mr. Walker's collection.
- ENCYRTUS *atricollis*, *Cat. Chalc. B. M.* p. 58. *Ratzeburg, Forst.*
Insekten, iii. t. 8, f. 13.
 b. Prussia. Presented by E. Doubleday, Esq.
- ENCYRTUS *ineptus*, *Cat. Chalc. B. M.* p. 59.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS *Marsus*, *Walker, Ent. Mag.* iv. 444.
 a. England. From Mr. Walker's collection.
- ENCYRTUS *Sipylus*, *Walker, Ent. Mag.* iv. 445.
 a. England. From Mr. Walker's collection.
- ENCYRTUS *Comara*, *Walker, Ent. Mag.* iv. 446.
 a. England. From Mr. Walker's collection.
- ENCYRTUS *Zarina*, *Haliday, Ent. Mag.* iv. 447.
 a. England. From Mr. Walker's collection.

- ENCYRTUS æneiventris, *Haliday, Ent. Mag.* iv. 447.
a. Isle of Bute. From Mr. Walker's collection.
- ENCYRTUS Jalysus, *Walker, Ent. Mag.* iv. 448.
a. England. From Mr. Walker's collection.
- ENCYRTUS Madyes, *Walker, Ent. Mag.* 448.
a. England. From Mr. Walker's collection.
- ENCYRTUS Chærilus, *Walker, Ent. Mag.* iv. 450.
a. England. From Mr. Walker's collection.
- ENCYRTUS Gellius, *Walker, Ent. Mag.* iv. 454.
a. England. From Mr. Walker's collection.
- ENCYRTUS Mattinus, *Walker, Ent. Mag.* iv. 455.
a. England. From Mr. Walker's collection.
- ENCYRTUS Machæras, *Walker, Ent. Mag.* iv. 460. *Ann. Nat. Hist.*
xiv. 185.
a. England. From Mr. Walker's collection.
- ENCYRTUS Ancharus, *Walker, Ent. Mag.* v. 36.
a. England. From Mr. Walker's collection.
- ENCYRTUS Atheas, *Walker, Ent. Mag.* v. 37.
a. England. From Mr. Walker's collection.
- ENCYRTUS Dius, *Walker, Ent. Mag.* v. 39.
a. England. From Mr. Walker's collection.
- ENCYRTUS Sosius, *Walker, Ent. Mag.* v. 39.
a. England. From Mr. Walker's collection.
- ENCYRTUS Corybas, *Walker, Ent. Mag.* v. 40.
a. England. From Mr. Walker's collection.
- ENCYRTUS Sosares, *Walker, Ent. Mag.* v. 41.
a. England. From Mr. Walker's collection.
- ENCYRTUS Pertiades, *Walker, Ent. Mag.* v. 42.
a. England. From Mr. Walker's collection.
- ENCYRTUS Thinxæus, *Walker, Ent. Mag.* v. 42.
a. England. From Mr. Walker's collection.
- ENCYRTUS Dercilus, *Walker, Ent. Mag.* v. 43.
a. England. From Mr. Walker's collection.
- ENCYRTUS Babas, *Walker, Ent. Mag.* v. 44.
a. England. From Mr. Walker's collection.
- ENCYRTUS Ariantes, *Walker, Ent. Mag.* v. 44.
a. England. From Mr. Walker's collection.
- ENCYRTUS Elbasus, *Walker, Ent. Mag.* v. 45.
a. England. From Mr. Walker's collection.

- ENCYRTUS Scaurus, *Walker, Ent. Mag.* v. 45.
 a. England. From Mr. Walker's collection.
- ENCYRTUS Jancirus, *Walker, Ent. Mag.* v. 46.
 a. England. From Mr. Walker's collection.
- ENCYRTUS Sitalces, *Walker, Ent. Mag.* v. 47.
 a. England. From Mr. Walker's collection.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS Tennes, *Walker, Ent. Mag.* v. 47.
 a. England. From Mr. Walker's collection.
- ENCYRTUS cyanocephalus, *Dalm. Kongl. Vet. Acad. Handl.* 1820.
 a. Sweden. From Mr. Walker's collection.
- ENCYRTUS cyanellus, *Dalm. Kongl. Vet. Acad. Handl.* 1820.
 a. Sweden. From Mr. Walker's collection.
- ENCYRTUS brevicornis, *Dalm. Kongl. Vet. Acad. Handl.* 1820.
 a. Sweden. From Mr. Walker's collection.
- ENCYRTUS melanacis, *Dalm. Kongl. Vet. Acad. Handl.* 1820.
 a. Sweden. From Mr. Walker's collection.
- ENCYRTUS lunatus, *Dalm. Kongl. Vet. Acad. Handl.* 1820.
 a. Sweden. From Mr. Walker's collection.
- ENCYRTUS longicornis, *Dalm. Kongl. Vet. Acad. Handl.* 1820.
 a. Sweden. From Mr. Walker's collection.
- ENCYRTUS Parus, *Walker, Ent. Mag.* v. 47.
 a. England. From Mr. Walker's collection.
- ENCYRTUS clavicornis, *Dalm. Kongl. Vet. Acad. Handl.* 1820. *Nees.*
Hym. Ich. aff. Mon. ii. 225.
 a. England. From Mr. Walker's collection.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENCYRTUS Eupales, *Walker, Ent. Mag.* v. 51.
 a. England. From Mr. Walker's collection.
- ENCYRTUS clavellatus, *Dalman, Kongl. Vet. Acad. Handl.* 1820.
 355, 41.
 a. Prussia. Presented by E. Doubleday, Esq.
 b. Prussia. From Mr. Walker's collection.
- ENCYRTUS Arceanus, *Walker, Ent. Mag.* v. 52.
 a. England. From Mr. Walker's collection.
- ENCYRTUS Tanais, *Walker, Ent. Mag.* v. 52.
 a. England. From Mr. Walker's collection.

ENCYRTUS Dahlbomii, *Westwood, Lond. and Edin. Phil. Mag. 3rd Series*, x. 63, 441. *Walker, Ent. Mag.* v. 53.

a. England. From Mr. Walker's collection.

ENCYRTUS hederaceus, *Westwood, Lond. and Edin. Phil. Mag. 3rd Series*, x. 63, 441.

a. England. From Mr. Walker's collection.

ENCYRTUS Erylus, *Walker, Ent. Mag.* v. 518.

a. England. From Mr. Walker's collection.

ENCYRTUS Dorens, *Walker, Ent. Mag.* v. 419.

a. England. From Mr. Walker's collection.

ENCYRTUS Amathus, *Walker, Ent. Mag.* v. 421.

a. England. From Mr. Walker's collection.

ENCYRTUS corniger, *Haliday, Walker, Ent. Mag.* v. 114.

a. France. From Mr. Walker's collection.

ENCYRTUS mirabilis, *Walker, Ent. Mag.* v. 114. Cerapterocerus mirabilis, *Westwood, Mag. Nat. Hist.* vi. 495. Encyrtus mirabilicornis, *Foerster, Mon. Pterom.* l. xliii. 2. Encyrtus poduroides, *Kollar.*

a. England. From Mr. Walker's collection.

ENCYRTUS fulvescens, *Walker, Ent. Mag.* v. 115. Ectroma fulvescens, *Westwood, Lond. and Edin. Phil. Mag. 3rd Series*, 111, 344.

a. England. From Mr. Walker's collection.

ENCYRTUS Gabestus, *Walker, Ent. Mag.* v. 116.

a. Ireland. From Mr. Walker's collection.

ENCYRTUS Barca, *Walker, Ent. Mag.* v. 116.

a. England. From Mr. Walker's collection.

ENCYRTUS Caris, *Walker, Ent. Mag.* v. 422.

a. England. From Mr. Walker's collection.

ENCYRTUS Nadius, *Walker, Ent. Mag.* v. 423.

a. England. From Mr. Walker's collection.

ENCYRTUS Camirus, *Walker, Ent. Mag.* v. 425.

a. England. From Mr. Walker's collection.

ENCYRTUS Telesto, *Walker, Ent. Mag.* v. 425.

a. England. From Mr. Walker's collection.

ENCYRTUS Meon, *Walker, Ent. Mag.* v. 426.

a. England. From Mr. Walker's collection.

ENCYRTUS Epona, *Walker, Ann. Nat. Hist.* xiv. 184.

a. England. From Mr. Walker's collection.

- ENCYRTUS Pyttalus, *Walker, Ann. Nat. Hist.* xiv. 184.
a. England. From Mr. Walker's collection.
- ENCYRTUS Nicippe, *Walker, Ann. Nat. Hist.* iv. 234.
a. England. From Mr. Walker's collection.
- ENCYRTUS Thyra, *Walker, Ent. Mag.* v. 427.
a. France. From Mr. Walker's collection.
- ENCYRTUS Obodas, *Walker, Ent. Mag.* v. 428.
a. France. From Mr. Walker's collection.
- ENCYRTUS Baleus, *Walker, Ent. Mag.* v. 428.
a. France. From Mr. Walker's collection.
- ENCYRTUS Cedrenus, *Walker, Ent. Mag.* v. 112.
a. South of France. From Mr. Walker's collection.
- ENCYRTUS Myrlea, *Walker, Ent. Mag.* v. 112.
a. South of France. From Mr. Walker's collection.
- ENCYRTUS Arene, *Walker, Ent. Mag.* v. 429.
a. South of France. From Mr. Walker's collection.
- ENCYRTUS Fadus, *Walker, Ent. Mag.* v. 429.
a. South of France. From Mr. Walker's collection.
- ENCYRTUS Sagillus, *Walker, Ent. Mag.* v. 430.
a. South of France. From Mr. Walker's collection.
- ENCYRTUS Gastron, *Walker, Ann. Nat. Hist.* xx. 21.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- ENCYRTUS Vectius, *Walker, Ann. Nat. Hist.* xx. 21.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- ENCYRTUS Reate, *Walker, Ann. Nat. Hist.* xx. 22.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- ENCYRTUS Cervius, *Walker, Ann. Nat. Hist.* xvii. 177.
a. Scotland. From Mr. Walker's collection.
- ENCYRTUS Meges, *Walker, Ann. Nat. Hist.* xvii. 178.
a. England. From Mr. Walker's collection.
- ENCYRTUS Præcia, *Walker, Ann. Nat. Hist.* xvii. 178.
a. England. From Mr. Walker's collection.
- ENCYRTUS Peticus, *Walker, Ann. Nat. Hist.* xvii. 179.
a. England. From Mr. Walker's collection.
- ENCYRTUS Metharma, *Walker, Ann. Nat. Hist.* xvii. 179.
a. England. From Mr. Walker's collection.
- ENCYRTUS Genutius, *Walker, Ann. Nat. Hist.* xvii. 180.
a. England. From Mr. Walker's collection.

- ENCYRTUS Mamertus, *Walker, Ann. Nat. Hist.* xvii. 180.
a. England. From Mr. Walker's collection.
- ENCYRTUS Alycæus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Molos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Phrosime, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Tylissos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Iophon, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Acratos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Hydramon, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Philotis, *Walker, MSS.* See Appendix.
a. Hammerfest, Finmark. From Mr. Walker's collection.
- ENCYRTUS Idmon, *Walker, MSS.* See Appendix.
a. Alten, Finmark. From Mr. Walker's collection.
- ENCYRTUS Alemon, *Walker, MSS.* See Appendix. *E. ænescens?*
Zetterstedt. Fauna Lapponica.
a. England. From Mr. Walker's collection.
- ENCYRTUS Genetyllis, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENCYRTUS Anthores, *Walker, MSS.* See Appendix.
a. Ireland. From Mr. Walker's collection.
- ENCYRTUS Idya, *Walker, MSS.* See Appendix.
a. Alten, Finmark. From Mr. Walker's collection.

MYINA, *Nees.*

- MYINA flava. *Aphelinus flavus, Walker, Mon. Chal.* i. 8. *Eulophus flavus, Nees. Hym. Ich aff. Mon.* ii. 167.
a. England. From Mr. Walker's collection.
- MYINA Argiope. *Aphelinus Argiope, Walker, Mon. Chal.* i. 9.
a. England. From Mr. Walker's collection.

MYINA Idæus. Aphelinus Idæus, *Walker, Mon. Chal.* i. 12.

a. England. From Mr. Walker's collection.

MYINA Acœtes. Aphelinus Acœtes, *Walker, Ann. Nat. Hist.* xiv. 185.

a. England. From Mr. Walker's collection.

PTEROPTRIX, *Westwood.*

PTEROPTRIX Hemera, *Walker, Mon. Chal.* i. 14.

a. England. From Mr. Walker's collection.

PTEROPTRIX Janias, *Walker, Mon. Chal.* i. 17.

a. England. From Mr. Walker's collection.

PTEROPTRIX Acestes, *Walker, Mon. Chal.* i. 19.

a. England. From Mr. Walker's collection.

PTEROPTRIX Thione, *Walker, Mon. Chal.* i. 19.

a. England. From Mr. Walker's collection.

ENTEDON, *Dalman.*

ENTEDON viridis. E. Prodice, *Walker, Cat. Chalc. B. M.* p. 63.

Elachestus viridis, *Nees, Hym. Ich. aff. Mon.* ii. 138, 2.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

ENTEDON Zanara, *Cat. Chalc. B. M.* p. 64.

b. Prussia. From Mr. Walker's collection.

ENTEDON cothurnatus. E. Argon, *Walker, Cat. Chalc. B. M.* p.

64. Elachestus cothurnatus, *Nees, Hym. Ich. aff. Mon.* ii. 141, 8.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

ENTEDON metallicus. E. Epigonus, *Walker, Cat. Chalc. B. M.*

Eulophus metallicus, *Nees, Hym. Ich. aff. Mon.* ii. 176, 34.

b. Prussia. Presented by E. Doubleday, Esq.

c. Prussia. From Mr. Walker's collection.

ENTEDON turcicus, *Walker, Mon. Chal.* i. 23. Eulophus turcicus,

Nees, Hym. Ich. aff. Mon. ii. 155, 5.

a. England. From Mr. Walker's collection.

ENTEDON Dimas, *Walker, Mon. Chal.* i. 24.

a. England. From Mr. Walker's collection.

ENTEDON Phænna, *Walker, Mon. Chal.* i. 25.

a. England. From Mr. Walker's collection.

ENTEDON Damastes. *Closterocerus Damastes*, *Walker, Ann. Nat. Hist.* xx. 22.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

ENTEDON Mento, *Walker, Mon. Chal.* i. 28.

a. England. From Mr. Walker's collection.

ENTEDON Tegar, *Walker, Mon. Chal.* i. 28.

a. England. From Mr. Walker's collection.

ENTEDON Coesius, *Walker, Mon. Chal.* i. 30.

a. England. From Mr. Walker's collection.

ENTEDON Alcetas. *Derostenus Alcetas*, *Walker, Ann. Nat. Hist.* xi. 31.

a. England. From Mr. Walker's collection.

ENTEDON Butus, *Walker, Mon. Chal.* i. 32.

a. England. From Mr. Walker's collection.

ENTEDON Imbrasmus, *Walker, Ann. Nat. Hist.* xx. 23.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

ENTEDON Proclea, *Walker, Mon. Chal.* i. 33.

a. England. From Mr. Walker's collection.

ENTEDON Nautes, *Walker, Mon. Chal.* i. 33.

a. England. From Mr. Walker's collection.

ENTEDON Nephereus, *Walker, Mon. Chal.* i. 36.

a. England. From Mr. Walker's collection.

ENTEDON Amasis, *Walker, Mon. Chal.* i. 37.

a. England. From Mr. Walker's collection.

EETEDON Cilla, *Walker, Mon. Chal.* i. 37.

a. England. From Mr. Walker's collection.

ENTEDON Idyia, *Walker, Mon. Chal.* i. 39.

a. England. From Mr. Walker's collection.

ENTEDON Thonis, *Walker, Mon. Chal.* i. 44.

a. England. From Mr. Walker's collection.

ENTEDON Parsodes, *Walker, Mon. Chal.* i. 45.

a. England. From Mr. Walker's collection.

ENTEDON Phryne, *Walker, Mon. Chal.* i. 46.

a. England. From Mr. Walker's collection.

ENTEDON Melænis, *Walker, Mon. Chal.* i. 47.

a. England. From Mr. Walker's collection.

ENTEDON Thersamon, *Walker, Mon. Chal.* i. 48.

a. England. From Mr. Walker's collection.

- ENTEDON Ocyalus, *Walker, Mon. Chal.* i. 56.
a. England. From Mr. Walker's collection.
- ENTEDON Pontinus, *Walker, Mon. Chal.* i. 60.
a. England. From Mr. Walker's collection.
- ENTEDON Leucippus, *Walker, Mon. Chal.* i. 61.
a. England. From Mr. Walker's collection.
- ENTEDON Abrota, *Walker, Mon. Chal.* i. 62.
a. England. From Mr. Walker's collection.
- ENTEDON Chilo, *Walker, Mon. Chal.* i. 63.
a. England. From Mr. Walker's collection.
- ENTEDON Alphenus, *Walker, Mon. Chal.* i. 64.
a. England. From Mr. Walker's collection.
- ENTEDON Afranius, *Walker, Mon. Chal.* i. 65.
a. England. From Mr. Walker's collection.
- ENTEDON Aso, *Walker, Mon. Chal.* i. 66.
a. England. From Mr. Walker's collection.
- ENTEDON Thoe, *Walker, Mon. Chal.* i. 68.
a. England. From Mr. Walker's collection.
- ENTEDON Charaxus, *Walker, Mon. Chal.* i. 68.
a. England. From Mr. Walker's collection.
- ENTEDON Tanis, *Walker, Mon. Chal.* i. 73.
a. England. From Mr. Walker's collection.
- ENTEDON Æsopus, *Walker, Mon. Chal.* i. 74.
a. England. From Mr. Walker's collection.
- ENTEDON Eropus, *Walker, Mon. Chal.* i. 75.
a. England. From Mr. Walker's collection.
- ENTEDON Coilus, *Walker, Mon. Chal.* i. 79.
a. England. From Mr. Walker's collection.
- ENTEDON Phruron, *Walker, Mon. Chal.* i. 87.
a. England. From Mr. Walker's collection.
- ENTEDON Epaphus, *Walker, Mon. Chal.* i. 89.
a. England. From Mr. Walker's collection.
- ENTEDON Rhesus, *Walker, Mon. Chal.* i. 88.
a. England. From Mr. Walker's collection.
- ENTEDON Marica, *Walker, Mon. Chal.* i. 88.
a. England. From Mr. Walker's collection.
- ENTEDON Phaola, *Walker, Mon. Chal.* i. 89.
a. England. From Mr. Walker's collection.

- ENTEDON Metius, *Walker, Mon. Chal.* i. 90.
 a. England. From Mr. Walker's collection.
- ENTEDON Clymene, *Walker, Mon. Chal.* i. 91.
 a. England. From Mr. Walker's collection.
- ENTEDON Charino, *Walker, Mon. Chal.* i. 96.
 a. England. From Mr. Walker's collection.
- ENTEDON Termerus, *Walker, Mon. Chal.* i. 96.
 a. England. From Mr. Walker's collection.
- ENTEDON Abdera, *Walker, Mon. Chal.* i. 98.
 a. England. From Mr. Walker's collection.
- ENTEDON Syma, *Walker, Mon. Chal.* i. 98.
 a. England. From Mr. Walker's collection.
- ENTEDON Ergias, *Walker, Mon. Chal.* i. 100.
 a. England. From Mr. Walker's collection.
- ENTEDON Methion. *Eulophus caniculus, Foerster, Mon. Pterom.*
 i. xl. 18.
 a. England. From Mr. Walker's collection.
 b. Prussia. Presented by E. Doubleday, Esq.
 c. Prussia. From Mr. Walker's collection.
- ENTEDON Beon, *Walker, Mon. Chal.* i. 107.
 a. England. From Mr. Walker's collection.
- ENTEDON Acantha, *Walker, Mon. Chal.* i. 108.
 a. England. From Mr. Walker's collection.
- ENTEDON Carne, *Walker, Mon. Chal.* i. 123.
 a. England. From Mr. Walker's collection.
- ENTEDON Vagellius, *Walker, Mon. Chal.* i. 123.
 a. England. From Mr. Walker's collection.
- ENTEDON Erginnus, *Walker, Mon. Chal.* i. 124.
 a. England. From Mr. Walker's collection.
- ENTEDON Hersilia, *Walker, Ann. Nat. Hist.* iv. 235.
 a. England. From Mr. Walker's collection.
- ENTEDON rutilans, *Haliday, Walker, Ann. Nat. Hist.* iv. 235.
 a. England. From Mr. Walker's collection.
- ENTEDON Acantha, *Walker, Mon. Chal.* i. 107.
 a. England. From Mr. Walker's collection.
- ENTEDON Epeus, *Walker, Mon. Chal.* i. 118.
 a. England. From Mr. Walker's collection.
- ENTEDON Alaspharus, *Walker, Mon. Chal.* i. 109.
 a. England. From Mr. Walker's collection.

- ENTEDON Linus, *Walker, Mon. Chal.* i. 119.
a. England. From Mr. Walker's collection.
- ENTEDON Fufius, *Walker, Ann. Nat. Hist.* xvii. 182.
a. England. From Mr. Walker's collection.
- ENTEDON Cœdicus, *Walker, Ann. Nat. Hist.* xvii. 183.
a. England. From Mr. Walker's collection.
- ENTEDON Nautias, *Walker, Ann. Nat. Hist.* xvii. 183.
a. England. From Mr. Walker's collection.
- ENTEDON Glabrio, *Walker, Ann. Nat. Hist.* xvii. 184.
a. England. From Mr. Walker's collection.
- ENTEDON Clinus. Horismenus Clinus, *Walker, Ann. Nat. Hist.*
xiv. 408.
a. Alten, Finmark. From Mr. Walker's collection.
- ENTEDON Isander, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Nurscia, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Sauros, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Axia, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Antaradus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Erxias, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Nænia, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Stennos, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Amadocus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Temena, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Metagenes, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Ergeteles, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- ENTEDON Adreus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

ENTEDON Ictetas, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

EPICLERUS, *Haliday.*

EPICLERUS Panyas. Entedon Panyas, *Walker, Mon. Chal.* i. 120.

a. England. From Mr. Walker's collection.

EPICLERUS Temenus. Entedon Temenus, *Walker, Mon. Chal.* i. 122.

e. England. From Mr. Walker's collection.

EUDERUS, *Haliday.*

EUDERUS Mithras, *Walker, Mon. Chal.* i. 93.

a. England. From Mr. Walker's collection.

EUDERUS Herillus, *Walker, Ann. Nat. Hist.* xx. 23.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

ELACHESTUS, *Spinola.*

ELACHESTUS Saon. Eulophus Saon, *Walker, Mon. Chal.* i. 175.

a. England. From Mr. Walker's collection.

ELACHESTUS Eurybates. Eulophus Eurybates, *Walker, Mon. Chal.* i. 178.

a. England. From Mr. Walker's collection.

ELACHESTUS Ticida. Eulophus Ticida, *Walker, Mon. Chal.* i. 179.

a. England. From Mr. Walker's collection.

ELACHESTUS Levana, *Walker, Ann. Nat. Hist.* xx. 27.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EULOPHUS, *Geoffroy.*

EULOPHUS ramicornis, *Cat. Chalc. B. M.* p. 69.

a. Prussia. Presented by E. Doubleday, Esq.

b. Prussia. From Mr. Walker's collection.

EULOPHUS Gonippus, *Cat. Chalc. B. M.* p. 69.

a. Prussia. From Mr. Walker's collection.

EULOPHUS gallarum, Nees, *Hym. Ich. aff. Mon.* ii. 170, 24. *E. Euedoreschus*, Walker, *Cat. Chalc. B. M.* p. 72. *Ichneumon gallarum*, Linn. *Syst. Nat.* ii. 939, 65. *Faun. Suec.* 16, 35. *Vill. Ent. Linn.* iii. 206, 232. *Fabr. Syst. Ent.* iii. *Ent. Syst.* ii. 186, 218. *Diplolepis Gallarum*, Fabr. *Syst. Piez.* 150, 8. *Spin. Ins. Lig. Fasc.* iv. 227. *Pteromalus Gallarum*, Latr. *Gen. Crust. et Ins.* iv. 31.

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

EULOPHUS Orsinus, Walker, *Mon. Chal.* i. 126.

a. England. From Mr. Walker's collection.

EULOPHUS Drupes, Walker, *Mon. Chal.* i. 127.

a. England. From Mr. Walker's collection.

EULOPHUS Anatole, Walker, *Mon. Chal.* i. 126.

a. England. From Mr. Walker's collection.

EULOPHUS Gobryas, Walker, *Ann. Nat. Hist.* xx. 26.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EULOPHUS Florus, Walker, *Mon. Chal.* i. 127.

a. England. From Mr. Walker's collection.

EULOPHUS Calavius, Walker, *Ann. Nat. Hist.* xx. 24.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EULOPHUS Æpulo, Walker, *Mon. Chal.* i. 141.

a. England. From Mr. Walker's collection.

EULOPHUS Iphinoe, Walker, *Ann. Nat. Hist.* xx. 25.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EULOPHUS Eucritus, Walker, *Mon. Chal.* i. 142.

a. England. From Mr. Walker's collection.

EULOPHUS Cyriades, Walker, *Ann. Nat. Hist.* xx. 25.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EULOPHUS Minio, Walker, *Ann. Nat. Hist.* xx. 25.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

EULOPHUS Leodamas, Walker, *Mon. Chal.* i. 130.

a. England. From Mr. Walker's collection.

EULOPHUS Thespius, Walker, *Mon. Chal.* i. 127.

a. England. From Mr. Walker's collection.

EULOPHUS Sancus, Walker, *Mon. Chal.* i. 138.

a. England. From Mr. Walker's collection.

EULOPHUS Carbo, Walker, *Mon. Chal.* i. 138.

a. England. From Mr. Walker's collection.

- EULOPHUS Pronoe, *Walker, Mon. Chal.* i. 140.
a. England. From Mr. Walker's collection.
- EULOPHUS Tidius, *Walker, Mon. Chal.* i. 146.
a. England. From Mr. Walker's collection.
- EULOPHUS Folia, *Walker, Mon. Chal.* i. 147.
a. England. From Mr. Walker's collection.
- EULOPHUS Mania, *Walker, Mon. Chal.* i. 154.
a. England. From Mr. Walker's collection.
- EULOPHUS Piscus, *Walker, Mon. Chal.* i. 155.
a. England. From Mr. Walker's collection.
- EULOPHUS Drusilla, *Walker, Mon. Chal.* i. 156.
a. England. From Mr. Walker's collection.
- EULOPHUS Euryalus, *Haliday, Trans. Ent. Soc.* iii.
a. Ireland. From Mr. Walker's collection.
- EULOPHUS Æpytus, *Walker, Mon. Chal.* i. 159.
a. England. From Mr. Walker's collection.
- EULOPHUS Docilis, *Walker, Mon. Chal.* i. 159.
a. England. From Mr. Walker's collection.
- EULOPHUS Prothenor, *Walker, Mon. Chal.* i. 180.
a. England. From Mr. Walker's collection.
- EULOPHUS Attis, *Walker, Mon. Chal.* i. 181.
a. England. From Mr. Walker's collection.
- EULOPHUS Sithon, *Walker, Mon. Chal.* i. 181.
a. England. From Mr. Walker's collection.
- EULOPHUS Croton, *Walker, Mon. Chal.* i. 182.
a. England. From Mr. Walker's collection.
- EULOPHUS Zeugma, *Walker, Mon. Chal.* i. 183.
a. England. From Mr. Walker's collection.
- EULOPHUS Metanira, *Walker, Mon. Chal.* i. 183.
a. England. From Mr. Walker's collection.
- EULOPHUS Cleopater, *Walker, Mon. Chal.* i. 184.
a. England. From Mr. Walker's collection.
- EULOPHUS Diyllus, *Walker, Mon. Chal.* i. 185.
a. England. From Mr. Walker's collection.
- EULOPHUS Minyas, *Walker, Ann. Nat. Hist.* xx. 25.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- EULOPHUS Anaxippus, *Walker, Ann. Nat. Hist.* xvii. 182.
a. England. From Mr. Walker's collection.

- EULOPHUS* Agathyllus, *Walker, Zoologist*. iv. 1458.
a. England. From Mr. Walker's collection.
- EULOPHUS* Idrieus, *Walker, Ann. Nat. Hist.* xiv. 409.
a. Alten, Finmark. From Mr. Walker's collection.
- EULOPHUS* Mamurius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- EULOPHUS* Rhamnius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- EULOPHUS* stygius, *Boheman, MSS.* See Appendix.
a. Sweden. From Mr. Walker's collection.
- EULOPHUS* Pythodorus, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- EULOPHUS* Acalles, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- EULOPHUS* Veturius, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.
- EULOPHUS* Scyllis, *Walker, MSS.* See Appendix.
a. England. From Mr. Walker's collection.

CIRROSPILUS, Westwood.

- CIRROSPILUS* Eunapius, *Walker, Ann. Nat. Hist.* xx. 27.
a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- CIRROSPILUS* flavovarius. *Eulophus arcuatus, Foerster, Mon. Pterom.* i. xxxix. 8. *E. 4-maculatus, Foerster, Mon. Pterom.* i. xxxix. 6.
a. Prussia. Presented by E. Doubleday, Esq.
b. Prussia. From Mr. Walker's collection.
- CIRROSPILUS* Pacuvius, *Walker, Ann. Nat. Hist.* i. 383.
a. England. From Mr. Walker's collection.
- CIRROSPILUS* Phorbas, *Walker, Ann. Nat. Hist.* i. 384.
a. England. From Mr. Walker's collection.
- CIRROSPILUS* Arsames, *Walker, Ann. Nat. Hist.* i. 384.
a. England. From Mr. Walker's collection.
- CIRROSPILUS* Orelia, *Walker, Ann. Nat. Hist.* i. 450.
a. England. From Mr. Walker's collection.
- CIRROSPILUS* Eudora, *Walker, Ann. Nat. Hist.* i. 452.
a. England. From Mr. Walker's collection.
- CIRROSPILUS* Procles, *Walker, Ann. Nat. Hist.* i. 453.
a. England. From Mr. Walker's collection.

CIRROSPILUS Abastor, *Walker, Ann. Nat. Hist.* i. 453.
a. England. From Mr. Walker's collection.

CIRROSPILUS Smilis, *Walker, Mon. Chal.* i. 317.
a. England. From Mr. Walker's collection.

CIRROSPILUS Myron, *Walker, Mon. Chal.* i. 294.
a. England. From Mr. Walker's collection.

CIRROSPILUS Deldon, *Walker, Mon. Chal.* i. 294.
a. England. From Mr. Walker's collection.

CIRROSPILUS Aratus, *Walker, Ann. Nat. Hist.* i. 453.
a. Ireland. From Mr. Walker's collection.

DIGLYPHUS, *Haliday.*

DIGLYPHUS Poppæa, *Haliday, MSS.* See Appendix.
a. Ireland. From Mr. Walker's collection.

DIGLYPHUS Aculeo, *Walker, MSS.* See Appendix.

TETRASTICHUS, *Haliday.*

TETRASTICHUS Evonymellæ, *Cat. Chalc. B. M.* p. 73.
b. Prussia. Presented by E. Doubleday, Esq.
c. Prussia. From Mr. Walker's collection.

TETRASTICHUS Armæus, *Cat. Chalc. B. M.* p. 74. Eulophus setiseries, *Foerster, Mon. Pterom.* i. xxxix. 2.
b. Prussia. Presented by E. Doubleday, Esq.
c. Prussia. From Mr. Walker's collection.

TETRASTICHUS miser. T. Attalus, *Walker, Cat. Chalc. B. M.* p. 74.
Eulophus miser, *Nees, Hym. Ich. aff. Mon.* ii. 173, 28.
c. Prussia. Presented by E. Doubleday, Esq.
d. Prussia. From Mr. Walker's collection.

TETRASTICHUS Prosymna, *Cat. Chalc. B. M.* p. 75. Eulophus crinicornis? *Perris, Ann. Soc. Ent. France.* ix.

TETRASTICHUS Quercens, *Cat. Chalc. B. M.* p. 76.
b. Prussia. Presented by E. Doubleday, Esq.
c. Prussia. From Mr. Walker's collection.

TETRASTICHUS Rapo, *Cat. Chalc. B. M.* Diplolepis Microgastri, *Bouché, Naturgesch. Ins.* i. 168, 61. Pteromalus Microgasteris, *Nees, Hym. Ich. aff. Mon.* ii. 423, 17. b.
b. Prussia. Presented by E. Doubleday, Esq.
c. Prussia. From Mr. Walker's collection.

TETRASTICHUS Nerio, *Cat. Chalc. B. M.* p. 77. Eulophus Ericæ?
Dufour, Ann. Soc. Ent. vii.

TETRASTICHUS Leucone, *Cat. Chalc. B. M.* p. 79. Eulophus longi-
 caudatus, *Foerster, Mon. Pterom.* i. xl. 19.

c. Prussia. Presented by E. Doubleday, Esq.

d. Prussia. From Mr. Walker's collection.

TETRASTICHUS Metra. Cirrospilus Metra, *Walker, Ann. Nat. Hist.*
 ii. 201.

a. England. From Mr. Walker's collection.

TETRASTICHUS Euedochus. Cirrospilus Euedochus, *Walker, Ann.*
Nat. Hist. ii. 204.

a. England. From Mr. Walker's collection.

TETRASTICHUS Cyrrihus. Cirrospilus Cyrrihus, *Walker, Ann. Nat.*
Hist. ii. 350.

a. England. From Mr. Walker's collection.

TETRASTICHUS æneus, *Haliday, MSS. Ann. Nat. Hist.* 1848.

a. Ireland. From Mr. Walker's collection.

TETRASTICHUS Mycerinus. Cirrospilus Mycerinus, *Walker, Ann.*
Nat. Hist. ii. 350.

a. England. From Mr. Walker's collection.

TETRASTICHUS Tachos. Cirrospilus Tachos, *Walker, Ann. Nat.*
Hist. ii. 352.

a. England. From Mr. Walker's collection.

TETRASTICHUS Tulis. Cirrospilus Tulis, *Walker, Ann. Nat. Hist.*
 ii. 354.

a. England. From Mr. Walker's collection.

TETRASTICHUS Murcia. Cirrospilus Murcia, *Walker, Ann. Nat.*
Hist. iii. 177.

a. England. From Mr. Walker's collection.

TETRASTICHUS Abantidas. Cirrospilus Abantidas, *Walker, Ann.*
Nat. Hist. iii. 179.

a. England. From Mr. Walker's collection.

TETRASTICHUS Euagoras. Cirrospilus Euagoras, *Walker, Ann.*
Nat. Hist. iii. 179.

a. England. From Mr. Walker's collection.

TETRASTICHUS Chares. Cirrospilus Chares, *Walker, Ann. Nat.*
Hist. iii. 180.

a. England. From Mr. Walker's collection.

- TETRASTICHUS Beroe. Cirrospilus Beroe, *Walker, Ann. Nat. Hist.* iii. 181.
a. England. From Mr. Walker's collection
- TETRASTICHUS Rhacius, *Walker, Ann. Nat. Hist.* iii. 181.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Æga. Cirrospilus Æga, *Walker, Ann. Nat. Hist.* iii. 181.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Anticlea. Cirrospilus Anticlea, *Walker, Ann. Nat. Hist.* iii. 182.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Sotades. Cirrospilus Sotades, *Walker, Ann. Nat. Hist.* iii. 417.
a. England. From Mr. Walker's collection
- TETRASTICHUS Anyta. Cirrospilus Anyta, *Walker, Ann. Nat. Hist.* iii. 417.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Zenocia. Cirrospilus Zenocia, *Walker, Ann. Nat. Hist.* iii. 418.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Zopyrus. Cirrospilus Zopyrus, *Walker, Ann. Nat. Hist.* iv. 29.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Arathis. Cirrospilus Arathis, *Walker, Ann. Nat. Hist.* iv. 29.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Rhode. Cirrospilus Rhode, *Walker, Ann. Nat. Hist.* iv. 29.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Clito. Cirrospilus Clito, *Walker, Ann. Nat. Hist.* iv. 30.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Rhosaces. Cirrospilus Rhosaces, *Walker, Mon. Chal.* i. 293.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Myron. Cirrospilus Myron, *Walker, Mon. Chal.* i. 294.
a. England. From Mr. Walker's collection.

- TETRASTICHUS Deldon. Cirrospilus Deldon, *Walker, Mon. Chal.*
i. 294.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Lycidas. Cirrospilus Lycidas, *Walker, Mon. Chal.*
i. 295.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Sinope. Cirrospilus Sinope, *Walker, Mon. Chal.*
i. 295.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Myrsus. Cirrospilus Myrsus, *Walker, Mon. Chal.*
i. 296.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Conon. Cirrospilus Conon, *Walker, Mon. Chal.*
i. 296.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Voranus. Cirrospilus Voranus, *Walker, Mon. Chal.*
i. 298.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Simo. Cirrospilus Simo, *Walker, Mon. Chal.* i.
298.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Plangon. Cirrospilus Plangon, *Walker, Mon. Chal.*
i. 298.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Vaccus. Cirrospilus Vaccus, *Walker, Mon. Chal.*
i. 299.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Oxathres. Cirrospilus Oxathres, *Walker, Mon.*
Chal. i. 299.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Eleuchia. Cirrospilus Eleuchia, *Walker, Mon. Chal.*
i. 301.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Dotus. Cirrospilus Dotus, *Walker, Mon. Chal.*
i. 301.
a. England. From Mr. Walker's collection.
- TETRASTICHUS Xeuxes. Cirrospilus Xeuxes, *Walker, Mon. Chal.*
i. 301.
a. England. From Mr. Walker's collection.

- TETRASTICHUS Ione. *Cirrospilus Ione*, *Walker, Mon. Chal.* i. 301.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Vacuna. *Cirrospilus Vacuna*, *Walker, Mon. Chal.* i. 305.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Oropus. *Cirrospilus Oropus*, *Walker, Mon. Chal.* i. 305.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Agrus. *Cirrospilus Agrus*, *Walker, Mon. Chal.* i. 306.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Tyrtæus. *Cirrospilus Tyrtæus*, *Walker, Mon. Chal.* i. 306.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Vicellius. *Cirrospilus Vicellius*, *Walker, Mon. Chal.* i. 307.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Epidius, *Walker, Ann. Nat. Hist.* xx. 28.
 a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.
- TETRASTICHUS Lyridice. *Cirrospilus Lyridice*, *Walker, Mon. Chal.* i. 307.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Gaus. *Cirrospilus Gaus*, *Walker, Mon. Chal.* i. 308.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Lachares. *Cirrospilus Lachares*, *Walker, Mon. Chal.* i. 309.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Apama. *Cirrospilus Apama*, *Walker, Mon. Chal.* i. 310.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Faucula. *Cirrospilus Faucula*, *Walker, Mon. Chal.* i. 310.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Racilla. *Cirrospilus Racilla*, *Walker, Mon. Chal.* i. 312.
 a. England. From Mr. Walker's collection.
- TETRASTICHUS Pamyles. *Cirrospilus Pamyles*, *Walker, Mon. Chal.* i. 313.
 a. England. From Mr. Walker's collection.

TETRASTICHUS Hypsistus. Cirrospilus Hypsistus, *Walker, Mon. Chal.* i. 316.

a. England. From Mr. Walker's collection.

TETRASTICHUS Smilis. Cirrospilus Smilis, *Walker, Mon. Chal.* i. 317.

a. England. From Mr. Walker's collection.

TETRASTICHUS Vincius. Cirrospilus Vincius, *Walker, Mon. Chal.* i. 317.

a. England. From Mr. Walker's collection.

TETRASTICHUS Eratus. Cirrospilus Eratus, *Walker, Mon. Chal.* i. 318.

a. England. From Mr. Walker's collection.

TETRASTICHUS Leocrates. Cirrospilus Leocrates, *Walker, Mon. Chal.* i. 319.

a. England. From Mr. Walker's collection.

TETRASTICHUS Totis. Cirrospilus Totis, *Walker, Mon. Chal.* i. 320.

a. England. From Mr. Walker's collection.

TETRASTICHUS Natras. Cirrospilus Natras, *Walker, Mon. Chal.* i. 320.

a. England. From Mr. Walker's collection.

TETRASTICHUS Hæmon, *Walker, Ann. Nat. Hist.* xx. 28.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

TETRASTICHUS Numeria. Cirrospilus Numeria, *Walker, Mon. Chal.* i. 321.

a. England. From Mr. Walker's collection.

TETRASTICHUS Anodaphus. Cirrospilus Anodaphus, *Walker, Mon. Chal.* i. 321.

a. England. From Mr. Walker's collection.

TETRASTICHUS Mutilia. Cirrospilus Mutilia, *Walker, Mon. Chal.* i. 322.

a. England. From Mr. Walker's collection.

TETRASTICHUS Gala, *Walker, Ann. Nat. Hist.* xx. 28.

a. St. John's Bluff, Florida. Presented by E. Doubleday, Esq.

TETRASTICHUS Trabea, *Walker, Mon. Chal.* i. 323.

a. England. From Mr. Walker's collection.

TETRASTICHUS Suevius. Cirrospilus Suevius, *Walker, Mon. Chal.* i. 323.

a. England. From Mr. Walker's collection.

TETRASTICHUS Rhæo. *Cirrospilus Rhæo*, *Walker, Ann. Nat. Hist.* iii. 417.

a. England. From Mr. Walker's collection.

TETRASTICHUS Teridæ. *Cirrospilus Teridæ*, *Walker, Ann. Nat. Hist.* iv. 236.

a. England. From Mr. Walker's collection.

TETRASTICHUS Brunchus. *Cirrospilus Brunchus*, *Walker, Ann. Nat. Hist.* iv. 236.

a. Scotland. From Mr. Walker's collection.

TETRASTICHUS Idothea, *Walker, Ann. Nat. Hist.* xiv. 409.

a. Alten, Finmark. From Mr. Walker's collection.

TETRASTICHUS Pausiris. *Cirrospilus Pausiris*, *Walker, Mon. Chal.* i. 327.

a. England. From Mr. Walker's collection.

TETRASTICHUS Marbis. *Cirrospilus Marbis*, *Walker, Mon. Chal.* i. 330.

a. England. From Mr. Walker's collection.

TETRASTICHUS Abydenus, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

TETRASTICHUS Bermius, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

TETRASTICHUS Axia, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

TETRASTICHUS Triarius, *Walker, MSS.* See Appendix.

a. England. From Mr. Walker's collection.

APPENDIX.

DESCRIPTONS OF NEW SPECIES

COMMUNICATED BY

FRANCIS WALKER, Esq., F.L.S., &c.

EURYTOMA Cestius. Fem. *Nigra, gracilis, genubus, tibiis apice tarsis, propedibusque rufis, alis limpidis.* Page 100.

The slender body of this species easily distinguishes it from all other described *Eurytomæ*; *E. atrum* comes nearest to it in this respect, but is very different in structure. Black, narrow, linear, convex: head and chest or thorax thickly punctured, not shining: head a little broader than the chest: feelers or antennæ subclavate, slender, shorter than the chest; club spindle-shaped, more than twice the length of the preceding joint: fore-chest or prothorax large; its breadth slightly exceeding its length: hind-chest or metathorax well developed, obconic, declining, having a large furrow down the middle: petiole short: abdomen spindle-shaped, smooth, shining, keeled beneath, a little longer and narrower than the chest; metapodeon and octoon of moderate size; ennaton shorter; deca-ton much longer; protelum, paratelum, and telum of moderate size: legs black; knees, feet or tarsi, and tips of shanks or tibiæ red; fore-legs nearly all red: wings white; veins tawny; ulna rather thick, about one-third of the length of the humerus; radius shorter than the ulna. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

Page 100² EURYTOMA Mucianus. Mas. *Nigra, genubus, tarsis propedibusque fulvis, alis limpidis.*

It resembles *E. platyptera*, but is distinguished from that species by its more nodose feelers. Black, convex: head and chest thickly punctured: feelers black, beset with whorls of hair, piceous

towards the tips, rather longer than the chest: breadth of the fore-chest about twice its length: petiole rather long: abdomen gibbous, very much shorter than the chest; its height is somewhat less than its length: legs black; knees and feet tawny, as are also the whole of the fore-legs except the base of the thighs: wings colourless; veins tawny; ulna not very broad, about one-fifth of the length of the humerus; radius longer than the ulna, a character wherein this *Eurytoma* differs from all the other British species, with the exception of *E. plumata* and *rufipes*. Length of the body 1 line; of the wings $1\frac{1}{2}$ line.

CALLIMOME Solinus. Mas. *Viridis, abdomine purpureo, antennis nigris, tibiis piceis, tarsis fulvis, alis subfuscis.*
Page 103.

Allied to *C. nigricornis*, but having a more slender body, with narrower and darker wings. Dark green, rather long and narrow: eyes dark red: feelers black, about half the length of the body, nearly filiform, moderately thick; first joint green: abdomen purple, rather narrower and shorter than the chest: hips, or coxæ, and thighs green; shanks piceous; knees, feet, and tips of the shanks tawny: wings narrow, and slightly tinged with brown; veins black; ulna nearly as long as the humerus; radius not more than one-third of the length of the ulna; cubitus less than one-half the length of the radius. Length of the body $1\frac{1}{2}$ lines; of the wings 3 lines.

ELATUS, *Walker*, (n. g.) Page 104.

This genus has a peculiar form, and somewhat resembles *Perilampus* and *Callimome*. *Male*. Body convex, arched: head and chest shagreened: head large, broader than the chest: feelers broad, subclavate, inserted in the middle of the front, much shorter than the chest; first joint long, slender; second cup-shaped; third and fourth very minute; fifth and following joints to the tenth short, broad, of nearly the same size, their breadth about twice their length; club conical, more than twice the length of the tenth joint: chest almost gibbous: fore-chest subquadrate, of moderate size; its breadth about twice its length: shield or scutum of the middle chest broad; sutures of the parapsides strongly marked; scutcheon or scutellum large, almost conical: hind-chest large, obconical, declining: petiole rather stout, shorter than the hind-chest: abdomen shorter and much narrower than the chest, smooth, shining, nearly elliptical, very convex, and forming a transverse ridge at the tip of

the metapodeon which extends over half the back; octoon also very large, and occupying the other half of the back, so that the following segments are not visible: ulna a little shorter than the humerus; radius about one-fourth of the length of the ulna; cubitus a little shorter than the radius, and hardly longer than in *Callimome* and *Perilampus*.

ELATUS Thenæ. *Cyaneo-viridis, abdomine æneo-viridi, antennis nigris, genubus tarsisque fuscis, tibiis apice fulvis, alis limpidis.* Page 104.

Bluish green: eyes red: feelers black; first joint green: abdomen tinged with bronze: legs green; tips of shanks tawny; knees and feet brown: wings colourless, of moderate size; veins brown; wing-brand or stigma, very small. Length of the body 1 line; of the wings $1\frac{3}{4}$ line.

TETRAMESA, *Walker*, (n. g.) Page 104.

Female. Convex: head and chest punctured, dull: head broader than the chest: feelers clavate, rather hairy, 13-jointed, as long as the chest; first joint long and slender; second nearly cup-shaped; third and fourth very small; fifth and following joints to the tenth slightly cup-shaped, successively increasing in breadth and decreasing in length; club 3-jointed, conical, full twice the length of the tenth joint: breadth of the fore-chest rather less than twice its length: sutures of the parapsides of the middle-chest strongly marked; axillæ large, separated from each other by a small interval: hind-chest obconical, declining, well developed: petiole short: abdomen nearly elliptical, smooth, shining, deeply keeled beneath, elevated at the tip, shorter but hardly narrower than the chest; metapodeon very large; octoon very short; ennaton long; decaton and following segments very short: wings rather narrow; ulna about one-fourth of the length of the humerus; radius shorter than the ulna; cubitus as long as the radius; wing-brand very small.

TETRAMESA *Iarbas*. Fem. *Nigra, genubus tarsisque rufis, alis subfuscis.* Page 104.

Black: knees and feet red; tips of the latter brown: wings slightly brown; veins piceous. Length of the body 1 line; of the wings $1\frac{3}{4}$ line.

GASTRANCISTRUS longicornis. Fem. *Nigro-viridis, antennis pedibusque piccis, genubus, tarsis tibiisque apice fulvis, alis limpidis*. Page 105.

Very dark green, almost black: head and chest very finely shagreened: head a little broader than the chest: eyes dark red: feelers clavate, slender, piceous, longer than the chest; first joint long and slender; second slightly cup-shaped; third and following joints to the seventh linear, nearly equal in size; club spindle-shaped, rather broader than the seventh joint and more than twice its length: chest oval, rather flat: fore-chest extremely short: shield of mid-chest broad; sutures of the parapsides indistinct; axillæ remote from each other; scutcheon nearly elliptical: hind-chest rather large, obconical, declining: petiole very short: abdomen very much compressed, knife-shaped, longer than the chest: sheaths of the oviduct about one-sixth of the length of the abdomen: legs piceous; knees, feet, and tips of the shanks tawny: wings colourless; veins tawny; ulna shorter than the humerus; radius more than one-third of the length of the ulna; cubitus shorter than the radius; wing-brand very small. Length of the body 1 line; of the wings $1\frac{1}{4}$ line.

GASTRANCISTRUS Amabæus. Fem. *Cyaneo-viridis, terebra abdominis dimidii longitudine, antennis nigris, pedibus flavis, alis limpidis*. Page 105.

Very nearly allied to *G. Dryas*, but the radius is somewhat shorter, and the colour of the legs is somewhat different. Bluish green: head and chest convex, finely shagreened: head as broad as the chest: feelers black, clavate, shorter than the chest; first joint stout, linear, tawny; second cup-shaped; third and fourth very small; fifth and following joints short, broad; club conical, about twice the length of the tenth joint: chest short, elliptical, very broad: fore-chest very short, not visible above: shield of middle-chest very large; sutures of the parapsides strongly marked, approaching each other behind; scutcheon rhomboidal, rather large; axillæ separated from each other by one-third of the breadth of the shield: hind-chest short, transverse, declining: petiole very short: abdomen spindle-shaped, greenish blue, compressed, smooth, shining, concave above, keeled beneath, a little longer than the chest, but hardly more than half its breadth, rather hairy towards the tip: sheaths of the oviduct black, hairy, full half the length of the abdomen: legs yellow; tips of the feet brown: wings colourless; forewings broad; veins brown; ulna but little shorter than the hume-

rus; radius not half the length of the ulna; cubitus much shorter than the radius; wing-brand of moderate size. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

GASTRANCISTRUS Accia. Fem. *Viridis, terebra brevissima, antennis nigris, tibiis fuscis, genubus tarsisque flavis, alis limpidis.* Page 105.

Allied to *G. annulipes*, but more slender. Bright green: head and chest convex, finely shagreened: head as broad as the chest: eyes red: feelers clavate, black, a little shorter than the chest; first joint long, slender; second cup-shaped; third and fourth very minute; fifth and following joints transverse, short, successively increasing in breadth; club ovate-conical, broader than the tenth joint and more than twice its length: chest elliptical; fore-chest very short; shield of the middle-chest large: sutures of the parapsides strongly marked, approaching each other; scutcheon rhomboidal: hind-chest obconical, declining, of moderate size: petiole very short: abdomen slender, spindle-shaped, smooth, shining, concave above, keeled beneath, longer and narrower than the chest; sheaths of the oviduct black, hairy, hardly one-fifth of the length of the abdomen: legs green; shanks brown; knees and feet yellow, tips of the latter brown: wings colourless; veins brown; ulna much shorter than the humerus; radius a little shorter than the ulna; cubitus much shorter than the radius; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{4}$ line.

GASTRANCISTRUS Hamillus. Fem. *Æneus, capite viridi, antennis fulvis, alis limpidis.* Page 105.

Allied to *G. autumnalis*. Brassy: head and chest convex, finely shagreened: head green, broader than the chest: feelers sub-clavate, tawny, as long as the chest; first joint long, slender, yellow; second cup-shaped; third and fourth very small; fifth and following joints closely joined together, successively increasing in breadth till the tenth; club spindle-shaped, pointed, not broader than the tenth joint, but more than twice its length, narrower than that of *G. autumnalis*: chest brassy, nearly elliptical; fore-chest short, just visible above; shield of the middle-chest rather broad; sutures of the parapsides strongly marked; axillæ separated from each other by at least one-third of the breadth of the shield; scutcheon nearly conical; hind-chest obconical, declining, of moderate size; petiole very short: abdomen elliptical, smooth, shining, concave above, slightly keeled beneath, narrower and a little longer

than the chest; metapodeon long; octoon and following segments shorter: legs bright yellow; tips of feet brown: wings colourless; fore-wings broader; veins yellow; ulna about half the length of the humerus; radius shorter than the ulna; cubitus still shorter; wing-brand small. Length of body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

GASTRANCISTRUS Geganjus. Mas. *Purpureus, abdomine basi viridi, antennis femoribusque fuscis, pedibus flavis, alis limpidis.* Page 105.

Purple: head and chest broad, convex, finely shagreened; head a little broader than the chest, inclining to bronze colour: eyes and eyelets or ocelli dark red: feelers 13-jointed, brown, clavate, much shorter than the chest; first joint spindle-shaped, long, yellow; second cup-shaped, pale brown; third and fourth very small; fifth and following joints transverse, slightly cup-shaped, successively decreasing in length; club oval, broader than the tenth joint, and more than twice its length: chest oval, thick, broader in front than behind; fore-chest very short, just visible above; shield of the middle-chest broad; sutures of the parapsides rather indistinct; axillæ separated by less than one-third of the breadth of the chest; scutcheon rhomboidal; hind-chest small, short-obconic, slightly declining, nearly smooth, with a brassy tinge, having a slight ridge along the middle; petiole very short: abdomen flat, linear, smooth, shining, green towards the base, hairy at the tip, a little longer than the chest, but not more than half its breadth; metapodeon long; octoon shorter; ennaton and following segments still shorter: legs yellow; thighs and shanks rather stout, the former brown: wings colourless; veins yellow; ulna not half the length of the humerus; radius a little longer than the ulna; cubitus as long as the ulna; wing-brand very small. Length of the body 1 line; of the wings $1\frac{1}{2}$ line.

GASTRANCISTRUS Fidenas. Mas. *Niger, antennis piceis, pedibus flavis, alis limpidis.* Page 105.

Black: head and chest black, convex, shagreened, dull; head large, broader than the chest: eyes piceous: eyelets red: feelers piceous, submoniliform, as long as the chest; first joint yellow, long and slender; second cup-shaped; third and fourth very minute; fifth and following joints to the tenth of moderate size, subquadrate; club ovate-conical, a little broader than the tenth joint and more than twice its length: chest nearly elliptical: fore-chest short, just visible above; shield of the middle-chest broad; sutures of the parapsides not very distinct, approaching each other behind; scutcheon

conical ; hind-chest large, obconical, declining : petiole short, stout : abdomen obclavate, flat, smooth, shining, a little shorter than the chest and not more than half its breadth ; metapodeon long ; octoon and following segments shorter : legs yellow : wings colourless ; veins pale tawny ; ulna much shorter than the humerus ; radius a little shorter than the ulna ; cubitus much shorter than the radius ; wing-brand very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

GASTRANCISTRUS Alectus. Mas. *Niger, abdomine nigro-æneo, petiolo fulvo, pedibus flavis, alis limpidis.* Page 105.

Head and chest convex, black, finely shagreened : head slightly tinged with green, broader than the chest : eyes and eyelets red : feelers brown, stout, filiform, longer than the chest ; first joint yellow, long, linear ; second cup-shaped ; third and fourth very small ; fifth and following joints to the ninth of moderate size, loosely joined together ; club spindle-shaped, more than twice the length of the ninth joint ; chest somewhat oval, thick, very convex, and having a slight brassy tinge ; fore-chest short, just visible above ; shield of the middle-chest broad ; sutures of the parapsides strongly marked ; axillæ separated by one-fourth of the breadth of the chest ; scutcheon large, rhomboidal, with a transverse suture before its tip ; hind-chest rather large, obconical, declining, with a transverse ridge along its middle : petiole short, tawny : abdomen dark bronze, smooth, shining, depressed, oval, green towards the base, shorter and narrower than the chest ; metapodeon rather long ; octoon and following segments shorter ; legs yellow ; feet paler, their tips tawny : wings colourless ; veins yellow ; ulna much shorter than the humerus ; radius as long as the ulna ; cubitus near one half the length of the radius ; wing-brand small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

GASTRANCISTRUS Hierocles. Mas. *Viridis, abdomine æneo basi subtus fulvo, antennis piceis, pedibus flavis, alis limpidis.* Page 105.

Green : head and chest convex, shagreened ; head a little broader than the chest ; eyes red ; feelers piceous, filiform, rather longer than the chest ; first joint long, slender ; second cup-shaped ; third and fourth very small ; fifth and following joints of nearly equal size : chest oval, rather short ; fore-chest very short ; shield of the mid-chest broad ; sutures of the parapsides very distinct ; axillæ separated by about one-third of the breadth of the chest ; scutcheon obconical ; hind-chest short ; petiole very short : abdomen dark

bronze, smooth, flat, nearly linear, tawny at the base beneath, narrower and a little longer than the chest; legs yellow; wings colourless; veins tawny; ulna much shorter than the humerus; radius as long as the ulna; cubitus much shorter than the radius; wing-brand of moderate size. Length of the body $\frac{1}{3}$ line; of the wings 1 line.

GASTRANCISTRUS Iriarte. Mas. *Viridis, abdomine æneo, antennis nigris, pedibus flavis, alis limpidis.* Page 105.

Bright green: head and chest convex, finely shagreened; head as broad as the chest, bright blue in front; feelers black, filiform, rather stout, about half the length of the body; first joint green, robust; second cup-shaped; third and fourth very small; fifth and following joints of nearly equal size: chest oval; fore-chest very short; shield of the middle-chest rather flat; sutures of the parapsides distinct; axillæ separated by nearly one half the breadth of the chest; scutcheon nearly obconical; hind-chest rather short, obconical, slightly declining: petiole very short: abdomen smooth, flat, nearly linear, dark bronze-colour, rather shorter and narrower than the chest; metapodeon rather long; octoon and following segments of moderate size: legs yellow; hips green: wings colourless; veins tawny; ulna much longer than the humerus; radius not one-fourth of the length of the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{4}$ line.

GASTRANCISTRUS Amnisos. Mas. *Æneo-viridis, abdomine purpureo basi subtus fulvo, antennis nigris, tibiis fulvis, tarsis flavis, alis subfuscis.* Page 105.

Brassy green: head and chest convex, shagreened; head very little broader than the chest; eyes dark red: feelers black, slightly moniliform, full as long as the chest; first joint green, slender, slightly curved; second cup-shaped; third and fourth very small; fifth and following joints of nearly equal size: chest spindle-shaped; fore-chest short; shield of the middle-chest rather flat, much narrower in front; sutures of the parapsides indistinct; axillæ approaching very near to each other; scutcheon somewhat convex, nearly elliptical; hind-chest rather long, obconical, slightly declining: abdomen slightly increasing in breadth from the base to the tip, flat, smooth, dark purple, tawny beneath at the base, a little shorter and much narrower than the chest; metapodeon rather longer than the following segments, which are of moderate and

nearly equal size; legs yellow; hips and thighs green; shanks tawny; wings slightly brown; veins brown; ulna much shorter than the humerus; radius a little longer than the ulna; cubitus rather long, about half the length of the radius; wing-brand small. Length of the body $1\frac{1}{4}$ line; of the wings $1\frac{3}{4}$ line.

GASTRANCISTRUS Pacilus. Fem. *Purpureus viridi varius, terebra abdominis trientis longitudine, antennis nigris, femoribus æneis tibiis fuscis, tarsis flavis, alis limpidis.* Page 105.

Purple, rather hairy: head and chest convex, finely shagreened; head not broader than the chest; crown tinged with green; eyes and eyelets dark red; feelers black, clavate, rather hairy, 13-jointed, a little shorter than the chest; first joint long, linear, brassy; second cup-shaped; third and fourth very small; fifth and following joints successively decreasing in length; club conical, broader than the tenth joint and more than twice its length; chest nearly elliptical, tinged with green; fore-chest very small, hardly seen above; shield of the middle chest rather long; sutures of the parapsides strongly marked; axillæ separated by full one-third the breadth of the chest; scutcheon nearly rhomboidal; hind-chest transverse, short, slightly declining; petiole very short; abdomen smooth, shining, rather hairy, nearly elliptical, slightly varied with green, depressed above, keeled beneath, a little narrower but not shorter than the chest; metapodeon long; ootoon and following segments shorter; sheaths of the oviduct black, hairy, nearly one-third of the length of the abdomen; hips and thighs brassy; shanks brown; knees and feet yellow, tips of the latter brown; wings colourless; veins brown; ulna about half the length of the humerus; radius shorter than the ulna; cubitus shorter than the radius, very slender; wing-brand rather large. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{4}$ line.

GASTRANCISTRUS Lælianus. Fem. *Viridi-cyaneus, abdominis disco æneo, oviductu brevi, antennis nigris, tarsis flavis, alis limpidis.* Page 105.

Greenish blue: head and chest convex, finely shagreened; head a little broader than the chest; eyes and eyelets red; feelers black, clavate, shorter than the chest; first joint green, long, slender; second cup-shaped; third and fourth very small; fifth and following joints short, transverse, successively decreasing in length; club conical, broader than the tenth joint and more than twice its

length : chest nearly oval ; fore-chest very short, hardly seen above ; shield of the middle-chest broad ; sutures of the parapsides strongly marked ; scutcheon nearly rhomboidal ; hind-chest obconic, declining, of moderate size : petiole very short : abdomen smooth, shining, bronze on the disk, depressed above, keeled beneath, decreasing in breadth from the base to the tip, narrower and much longer than the chest ; sheaths of the oviduct black, hairy, nearly one-fourth of the length of the abdomen : legs green ; knees yellow ; feet brown : wings very slightly tinged with brown ; veins brown ; ulna half the length of the humerus ; radius as long as the ulna ; cubitus much shorter than the radius ; wing-brand of moderate size. Length of the body $\frac{3}{4}$ line ; of the wings 1 line.

PSILONOTUS Achæus. Fem. *Aureo-viridis, antennis tibiisque piceis, tarsis flavis, genubus et protarsis fulvis, alis limpidis.*
Page 105.

Head and chest bright golden green, shining, convex, finely shagreened : head as broad as the chest ; front large, flat, produced into a small obtuse tooth between the feelers ; the latter are inserted by the mouth, piceous, subclavate, very slender, longer than the chest ; first joint long, slender, tawny, very slightly curved ; second long cup-shaped ; third and fourth small, but more distinct than is usual in this family ; fifth and following joints subquadrate, successively decreasing in length ; club spindle-shaped, a little broader than the ninth joint and more than twice its length ; chest elliptical ; fore-chest very short ; shield of the middle-chest broad, flat ; sutures of the parapsides indistinct ; axillæ large, separated by nearly one-third of the breadth of the chest ; scutcheon flat, nearly rhomboidal ; hind-chest short, declining ; petiole very short : abdomen lanceolate, much compressed, smooth, shining, concave above, keeled beneath, as long as the chest, but not more than one-third of its breadth : hips and thighs green ; knees and fore feet tawny ; shanks piceous, their tips and the four hinder feet yellow : wings colourless ; veins brown ; ulna full half the length of the humerus ; radius shorter than the ulna ; cubitus shorter than the radius ; wing-brand small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

MACROMESUS. Page 106.

Female. Head and chest convex, finely shagreened ; head as broad as the chest ; feelers 14-jointed, subclavate, pubescent ; first joint long, slender ; second very long, cup-shaped ; third and fourth very short ; fifth and following joints long, linear ; club long,

conical, more than twice the length of the eleventh joint: chest elliptical; fore-chest very short, hardly seen above; shield of the middle-chest rather long; sutures of the parapsides strongly marked; axillæ separated by nearly one half the breadth of the chest; scutcheon long, nearly rhomboidal: hind-chest short, transverse, declining: petiole very short: abdomen spindle-shaped, smooth, shining, depressed above, hardly keeled beneath: legs long; middle shanks twice the length of the fore-shanks and a little longer than the hind-shanks; middle-feet a little longer than the other feet.

MACROMESUS Amphiretus. Fem. *Æneus cyaneo viridi purpureoque varius, antennis piceis, pedibus fulvis, alis subfuscis.*
Page 106.

Body rather slender: head and chest bronze, slightly varied with green: feelers very slender, piceous, a little longer than the chest: first joint reddish yellow: front of the head varied with purple, blue and green: abdomen purple-bronze, slightly tinged with green at the base, rather hairy towards the tip, longer and a little broader than the chest: legs reddish yellow; tips of the feet brown; wings very slightly tinged with brown; veins brown; ulna rather more than half the length of the humerus; radius much shorter than the ulna; cubitus a little shorter than the radius; wing-brand of moderate size. Length of the body $1\frac{1}{2}$ line; of the wings $2\frac{1}{2}$ line. Allied to Merostenus and the Eupelmidæ?

PIRENE Scylax. Mas. *Viridis, antennis nigris, genubus, tarsis tibisque apice fulvis, tibiis propedibusque piceis, alis limpidis.* Page 106.

Dark green: head broader than the chest: eyes red: feelers black, clavate, shorter than the chest; first joint much dilated; second cup-shaped; third and following joints very small; club large, oval: abdomen compressed, nearly as long as the chest: hips and thighs dark green; knees, feet, and tips of shanks tawny; shanks and fore-feet piceous: wings colourless; veins piceous; ulna nearly as long as the humerus; radius not more than one-fourth of the length of the ulna; cubitus a little shorter than the radius; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

ORMOCERUS Sabbas. Fem. *Cyaneo-viridis, antennis fuscis basi viridibus, tibiis viridi-piceis, tarsis flavis, alis limpidis.*
Page 106.

Bright bluish green: head and chest convex, finely shagreened;

head a little broader than the chest: eyes and eyelets red: feelers brown, clavate, rather slender, a little longer than the chest; first joint green, long, slender; second cup-shaped, green; third and fourth very small; fifth and following joints successively, but very slightly decreasing in length; club elliptical, broader than the tenth joint and more than twice its length: chest oval, broader in front than behind; fore-chest very short, just seen above; shield of the middle-chest rather long; sutures of the parapsides rather strongly marked; axillæ separated by rather less than one-fourth of the breadth of the chest; scutcheon nearly rhomboidal, of moderate size, having a transverse suture near the tip; hind-chest short, transverse, declining; petiole very short: abdomen spindle-shaped, smooth, shining, concave above, slightly keeled beneath, longer and much narrower than the chest; disk slightly tinged with bronze; hips and thighs green; shanks piceous, tinged with green; knees and feet yellow, tips of the latter piceous; fore-feet brown: wings colourless; veins brown; ulna full half the length of the humerus; radius much longer than the ulna; cubitus much shorter than the ulna; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

ORMOCERUS Aletes. Fem. *Aureo-viridis, abdomine cyaneo-viridi, antennis piceis, tibiis fulvis, pedibus flavis, alis limpidis.* Page 107.

Body thick and broad: head and chest golden-green, convex, finely shagreened: head very large, broader than the chest: eyes and eyelets red: feelers 12-jointed, piceous, clavate, stout, short, not more than half the length of the chest; first joint long, linear, tawny, piceous at the tip; second cup-shaped; third and fourth very small; fifth and following joints short, transverse, successively decreasing in length; club elliptical, a little broader than the ninth joint and more than twice its length: chest deep, very robust; fore-chest very short; shield of the middle-chest very broad; sutures of the parapsides very indistinct; axillæ separated by one-fourth of the breadth of the chest; scutcheon truncate-conic, having a very slight trace of a transverse suture near its tip; hind-chest of moderate size, obconical, declining, with a ridge along the middle and one on each side by which it is divided into two compartments; petiole extremely short: abdomen bluish green, smooth, shining, concave above, very deeply keeled beneath, narrower but hardly shorter than the chest; the disk is slightly tinged with bronze; metapodeon rather large; octoon and following segments of moderate size: legs green; shanks tawny; knees and tips of shanks yellow; feet yellow, their tips piceous: wings colourless; veins brown; ulna

rather more than half the length of the humerus; radius shorter than the ulna; cubitus shorter than the radius; wing-brand small. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

CYRTOGASTER PÆSOS. Fem. *Viridis, abdomine æneo, antennis nigris, pedibus luteis, metatarsis flavis, alis limpidis.*
Page 108.

Head and chest convex, finely shagreened: head bluish green, a little broader than the chest: eyes and eyelets red: feelers black, clavate, a little shorter than the chest; first joint green; second cup-shaped; third and fourth very small; fifth and following joints short, and successively decreasing in length; club elliptical, broader than the tenth joint and more than twice its length: chest green, nearly oval, not very convex; fore-chest very short; shield of the middle-chest rather short; sutures of the parapsides very indistinct; axillæ separated by rather more than one-fourth of the breadth of the chest; scutcheon nearly rhomboidal; hind-chest large, obconical, declining, rather roughly shagreened: petiole moderately long: abdomen dark bronze, oval, smooth, shining, convex above, keeled beneath, a little shorter and broader than the chest; metapodeon and octoon rather large; ennaton and following segments short: legs luteous; hips green; middle-feet and hind-feet yellow, their tips piceous: wings colourless; veins tawny; ulna rather more than half the length of the humerus; radius much shorter than the ulna; cubitus a little shorter than the radius; wing-brand small. Length of the body $\frac{2}{3}$ line; of the wings $1\frac{1}{3}$ line.

TITYROS, Walker. Page 108.

Male. Head and chest convex, finely shagreened: head very large and thick, much broader than the chest: feelers clavate, as long as the chest; first joint nearly linear, very long and slender, nearly as long as all the following joints; second cup-shaped; third and fourth joints very small: fifth and following joints very short; club elliptical, much broader than the tenth joint and more than twice its length: chest short-oval, deep, broad; fore-chest very short; shield of the middle-chest large; sutures of the parapsides indistinct; scutcheon nearly conical; hind-chest large, obconical, declining, nearly smooth: petiole short: abdomen nearly conical, depressed, smooth, shining, much shorter than the chest and not more than half its breadth; metapodeon occupying most of the back; octoon and following segments very short: ulna not more than half the length of the humerus; radius a little shorter than the ulna; cubitus much shorter than the radius; wing-brand very small.

TITYROS Poreia. Mas. *Viridis, antennis fuscis basi viridibus, tibiis tarsis que flavis, illis fulvo cinctis, protibiis et protarsis luteis, alis limpidis.* Page 108.

Green : eyes and eyelets red : feelers brown ; first joint green, yellow at the base ; second piceous, tinged with green : hips and thighs green ; tips of the latter yellow ; trochanters tawny ; shanks and feet of the four hinder legs yellow, the former traversed by a broad tawny band, tips of the latter piceous ; fore-shanks and fore-feet luteous, tips of the latter brown : wings colourless ; veins tawny ; humerus brown. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{4}$ line.

SPHEGIGASTER Agriope. Mas. *Cyaneo-viridis, antennis nigris, pedibus luteis, alis limpidis.* Page 108.

Bright bluish green, convex : head and chest finely shagreened : head blue, much broader than the chest : eyes red : feelers black, filiform, slender, much longer than the chest ; first joint yellow, slender, linear ; second cup-shaped ; third and fourth very small ; fifth and following joints long, linear ; club spindle-shaped, much longer than the preceding joint : chest nearly elliptical : fore-chest short : shield of the mid-chest much narrower in front ; sutures of the parapsides distinct ; axillæ large, meeting on the back ; scutcheon nearly rhomboidal : hind-chest long, obconical, declining : petiole slender, cylindrical, nearly half the length of the abdomen : abdomen oval, smooth, a little narrower and very much shorter than the chest ; metapodeon concave on the disk, occupying nearly half the back ; octoon and ennaton of moderate size ; decaton and following segments short : legs luteous ; hips green : wings colourless ; veins tawny ; ulna very much shorter than the humerus ; radius longer than the ulna ; cubitus about half the length of the ulna ; wing-brand small. Length of the body $1-1\frac{1}{4}$ line ; of the wings $2-2\frac{1}{4}$ lines.

SPHEGIGASTER Lasthenes. Fem. *Viridis, abdomine purpureo-æneo basi cyaneo, antennis nigris, pedibus luteis, femoribus viridibus tarsis flavis, alis limpidis.* Page 108.

Dark green, convex : head and chest shagreened : head much broader than the chest : eyes dark red : feelers black, clavate, nearly as long as the chest ; first joint slender, linear, luteous, black at the tip ; second cup-shaped ; third and fourth very small ; fifth and following joints to the tenth successively increasing in

breadth; club conical at the tip, a little broader than the tenth joint, and more than twice its length: chest elliptical: fore-chest very short: shield of the mid-chest much narrower in front; sutures of the parapsides indistinct; axillæ large, approaching near to each other; scutcheon somewhat rhomboidal: hind-chest long, obconical, declining, with a rim on each side: petiole stout, cylindrical, shagreened, about one-sixth of the length of the abdomen: abdomen smooth, nearly elliptical, purplish bronze on the disk, blue towards the base, deeply keeled beneath, nearly as broad as the chest but very much shorter; metapodeon long, occupying nearly half the back; octoon of moderate size; ennaton and following segments very short: legs luteous; hips and the greater part of the thighs green; middle-feet and hind-feet yellow, their tips and those of the fore-feet piceous: wings colourless; veins piceous; ulna little more than half the length of the humerus; radius very little longer than the ulna; cubitus more than half the length of the ulna; wing-brand very small. Length of the body 1 line; of the wings 2 lines.

LAMPROTATUS Mallius. Mas. *Æneo-viridis, antennis nigris, pedibus fulvis, femoribus viridi cinctis, metatibiis apice tarsisque piceis, alis fuscis.* Page 111.

Coppery green, convex: head and chest very finely shagreened, appearing almost smooth: head a little broader than the chest: eyes and eyelets red: feelers black, filiform, longer than the chest; first joint long, slender, green; second cup-shaped; third and fourth very small; fifth and following joints long, linear, of nearly equal length; club spindle-shaped, much longer than the tenth joint: chest oval: fore-chest transverse, narrower in front; its length above one-fourth of its breadth; the hind border concave: shield of the mid-chest rather long; parapsides much developed, their sutures very strongly marked; axillæ rather large, parted by about one-fifth of the breadth of the chest; scutcheon subrhomboidal, with a transverse suture near its tip, a longitudinal furrow on each side, and a slight trace of a furrow proceeding from the middle of the base towards the disk: hind-chest rather large, obconical, declining; its fore-shield distinct: petiole about one-seventh of the length of the abdomen: abdomen oval, smooth, a little narrower and much shorter than the chest; metapodeon occupying nearly half of the back, its disk concave from the base till near the tip: octoon about half the length of the metapodeon; ennaton much shorter; decaton and following segments still shorter: legs tawny; hips green; feet piceous, the joints paler towards the base; fore-thighs and hind-thighs traversed by a broad green band; tips

of hind-shanks piceous: wings brown; veins piceous; humerus hardly twice the length of the ulna; radius very much longer than the ulna; cubitus about half the length of the ulna; wing-brand large. Length of the body $1\frac{1}{2}$ line; of the wings $2\frac{1}{2}$ lines.

LAMPROTATUS Zipoetes. Fem. *Cyaneo-viridis, abdomine æneo-viridi, antennis nigris, tibiis piceis, tarsis flavis, alis subfuscis.* Page 111.

Convex: head and chest finely shagreened: head bluish green, much broader than the chest: eyes and eyelets red: feelers black, clavate, much shorter than the chest; first joint long, linear, green; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest elliptical, greenish blue, tinged with purple; fore-chest transverse; its length above about one-sixth of its breadth: shield of the mid-chest long; parapsides well developed, their sutures strongly marked; axillæ parted by about one-fifth of the breadth of the chest; scutcheon subrhomboidal, with a transverse suture near its tip: hind-chest rather large, obconical, declining, with a slight furrow along the middle and a rim on each side: petiole about one-seventh of the length of the abdomen: abdomen brassy green, elliptical, smooth, keeled beneath, a little narrower and much shorter than the chest; metapodeon rather more than one-third of its length; octoon about half the length of the metapodeon; ennaton as long as the octoon; decaton a little shorter; the following segments very short: hips and thighs green; shanks piceous; knees, feet, and tips of shanks yellow; tips of feet brown; fore-shanks and fore-feet tawny: wings slightly tinged with brown; veins brown; ulna hardly half the length of the humerus; radius much longer than the ulna; cubitus more than half the length of the ulna; wing-brand piceous, rather large. Length of the body 1 line; of the wings 2 lines.

LAMPROTATUS Œbares. Fem. *Cyaneo-viridis, antennis nigris, tibiis tarsisque fulvis apice piceis, alis limpidis.* Page 111.

Bluish green, convex: head and chest finely shagreened: head a little broader than the chest: eyes and eyelets red: feelers black, clavate, a little longer than the chest; first joint green, long, slender; second cup-shaped; third and fourth very small; fifth and following joints to the tenth successively decreasing in length;

club long-oval, broader than the tenth joint and more than twice its length : chest oval : fore-chest very short ; its length above hardly one-fourth of its breadth : sutures of the parapsides strongly marked ; axillæ approaching near to each other ; scutcheon subrhomboidal, with a transverse suture near its tip : hind-chest large, obconical, declining : petiole short : abdomen oval, smooth, keeled beneath, a little broader and much shorter than the chest ; metapodeon occupying about half of the back ; octoon and following segments of moderate size : legs tawny ; hips and thighs green, tips of the latter tawny ; shanks and feet of four hinder legs with piceous tips : wings colourless ; veins brown ; ulna about half the length of the humerus ; radius very much longer than the ulna ; cubitus full half the length of the ulna ; wing-brand small. Length of the body $\frac{2}{3}$ line ; of the wings $1\frac{1}{2}$ line.

LAMPROTATUS Phylander. *Æneo-viridis, pedibus piceis, tarsis flavis, protarsis fulvis, alis limpidis.* Page 111.

Head wanting : chest coppery green, convex, finely shagreened : fore-chest rather short ; its length about one-third of its breadth ; sutures of the parapsides indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal ; hind-chest obconical, declining, with a slight ridge along the middle : petiole moderately long : abdomen deep, oval, smooth, convex, brassy green, slightly keeled beneath, much shorter but not narrower than the chest ; metapodeon occupying more than one-third of the back ; octoon and ennaton of moderate size ; decaton and following segments very short : legs piceous ; hips green ; knees and feet yellow, tips of the latter brown ; fore-shanks and fore-feet tawny : wings colourless ; veins pale brown ; ulna much shorter than the humerus ; radius nearly as long as the ulna ; cubitus not half the length of the radius ; wing-brand very small. Length of the body $\frac{2}{3}$ line ; of the wings $1\frac{1}{2}$ line.

LAMPROTATUS Acarnas. Fem. *Viridis, abdominis disco purpureo, antennis nigris, pedibus flavis, alis limpidis.* Page 111.

Green, convex : head and chest finely shagreened : head bluish green, a little broader than the chest : eyes and eyelets red : feelers black, clavate, a little shorter than the chest ; first joint yellow, long, linear, piceous at the tip ; second long cup-shaped ; third and fourth very small ; fifth and the following joints to the tenth suc-

cessively decreasing in length; club long-conical, broader than the tenth joint and more than twice its length: chest oval: fore-chest very short: shield of the mid-chest broad; sutures of the parapsides indistinct; axillæ parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal, having a slight transverse suture near its tip: hind-chest of moderate size, obconical, declining: petiole rather short: abdomen elliptical, smooth, almost flat above, slightly keeled beneath, much narrower and shorter than the chest; metapodeon occupying about half the back; octoon dark purple, large; ennaton and following segments extremely short: legs yellow; hips green; four hinder-feet pale yellow with brown tips: wings colourless; veins tawny; ulna about half the length of the humerus; radius much longer than the ulna; cubitus more than half the length of the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $\frac{1}{2}$ line.

LAMPROTATUS Acilius. Mas. *Viridis, antennis fuscis, pedibus flavis, alis limpidis.* Page 111.

Bright green, convex, narrow: head and chest finely shagreened: head much broader than the chest: eyes and eyelets red: feelers pale brown, filiform, rather shorter than the chest; first joint green, long, slender; second long-cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest spindle-shaped: fore-chest very short, much narrower than the middle-chest, with the hind-border nearly straight; its length above about one-fourth of its breadth: shield of the middle-chest rather long; sutures of the parapsides very distinct; axillæ parted by about one-fourth of the breadth of the chest; scutcheon subrhombiform: hind-chest rather long, obconical, declining, with a ridge along the middle and a rim on each side: petiole short: abdomen oval, smooth, hardly half the length of the chest; metapodeon occupying half of the back; octoon short; ennaton and following segments very short: legs yellow; hips green; four hinder-feet pale yellow with piceous tips: wings colourless, rather narrow; veins yellow; ulna about half the length of the humerus; radius as long as the ulna; cubitus more than half the length of the ulna; wing-brand extremely small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

LAMPROTATUS Amulius. Fem. *Viridi-æneus, antennis nigris, pedibus luteis, femoribus basi viridibus, tarsis flavis, alis limpidis.* Page 111.

Greenish blue, convex: head and chest shagreened: head not

broader than the chest: eyes and eyelets red: feelers black, subclavate, a little longer than the chest; first joint green; second cup-shaped; third and fourth very small; fifth and following joints to the tenth successively decreasing in length; club spindle-shaped; much more than twice the length of the tenth joint: chest oval: fore-chest very short, narrow in front, slightly concave behind; its length above not more than one-sixth of its breadth: shield of the middle-chest broad; sutures of the parapsides very distinct; axillæ rather large, parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a transverse suture near its tip: hind-chest large, obconical, declining: petiole very short: abdomen short-elliptical, smooth, keeled beneath, a little narrower and very much shorter than the chest; metapodeon occupying full half of the back, concave at the base; octoon and following segments short: legs luteous; hips and the lower part of the thighs green; four hinder-feet yellow with brown tips: wings colourless; veins brown; ulna about half the length of the humerus; radius very much longer than the ulna; cubitus full half the length of the ulna; wing-brand rather large. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

LAMPROTATUS Labaris. Mas. *Viridis cyaneo varius, abdominis disco aeneo, antennis nigris, pedibus fulvis, tarsis flavis, profemoribus piceo vittatis, alis limpidis.* Page 111.

Bright green: head and fore-chest bluish green: head and chest convex, shagreened: head a little broader than the chest: eyes and eyelets red: feelers black, filiform, longer than the chest; first joint green; second cup-shaped; third and fourth very small; fifth and following joints successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest oval: fore-chest short, narrow in front, concave behind; its breadth above more than twice its length: sutures of the parapsides very distinct; axillæ large, approaching each other; scutcheon subrhomboidal, with a transverse suture near the tip: hind-chest large, obconical, declining: petiole short: abdomen smooth, flat, nearly linear, slightly contracted at the base, narrower and very much shorter than the chest; disk bronze colour; metapodeon large; octoon and following segments short: legs tawny; hips green; each fore-thigh with a piceous line along its whole length; knees and feet of four hinder-legs yellow, tips of the latter brown: wings colourless; veins piceous; humerus much longer than the ulna; radius hardly longer than the ulna; cubitus full half the length of the ulna; wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{4}$ lines.

LAMPROTATUS Bolgius. Fem. *Æneus, abdomine viridi vario, antennis nigris, pedibus fulvis, tarsis flavis, metafemoribus basi piceis, alis limpidis.* Page 111.

Brassy, convex: head and chest shagreened: head tinged with green, hardly broader than the chest: eyes and eyelets dark red: feelers black, pubescent, subclavate, not longer than the chest; first joint long, slender, green; second cup-shaped; third and fourth very small; fifth and following joints to the tenth successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest oval: fore-chest short; its breadth above more than twice its length: sutures of the parapsides distinct; axillæ rather large, parted by about one-fourth of the breadth of the chest; scutcheon nearly conical: hind-chest large, obconical, declining, with a ridge along the back: petiole short: abdomen oval, smooth, tinged with green, keeled beneath, much shorter but not narrower than the chest; metapodeon occupying about one-third of the back; octoon and three following segments of moderate size; paratelum and telum very short: legs tawny; hips brassy; base of the hind-thighs piceous; middle-feet and hind-feet yellow with brown tips: wings colourless; veins tawny; ulna rather more than half the length of the humerus; radius much longer than the ulna; cubitus more than half the length of the ulna; wing-brand small. Length of the body $\frac{3}{4}$ lines; of the wings $1\frac{1}{2}$ line,

LAMPROTATUS Methymna. Mas. et Fem. *Viridis, aneo aut cyaneo varius, antennis nigris, pedibus piceis, alis fuscis.* Page 111.

Male. Dark green, with a slight bluish tinge: head and chest convex, very finely shagreened, appearing almost smooth: head large, broader than the chest: feelers black, filiform, pubescent, much longer than the chest; first joint long, slender, green; second cup-shaped; third and fourth very small; fifth and following joints to the tenth long, linear, of equal length; club spindle-shaped, much longer than the tenth joint: chest oval, well developed: fore-chest short; its breadth above about four times its length; sutures of the parapsides very strongly developed; axillæ large, and nearly meeting on the back; scutcheon subrhomboidal, with a transverse suture near its tip: hind-chest large, obconical, declining; its fore-shield transverse, short, distinct: petiole short: abdomen brassy green, smooth, depressed, increasing in breadth from the base to the tip, shorter and much narrower than the chest;

metapodeon occupying much more than half of the back; octoon and ennaton of moderate size; decaton and following segments very short: legs piceous; hips green; knees and fore-shanks pale piceous: wings brown, long and large; veins dark piceous; ulna about half the length of the humerus; radius nearly twice the length of the ulna; cubitus full half the length of the ulna; wing-brand large. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{2}$ lines.

Var. *a*. Head and chest green.

Var. *β*. Head and chest brassy green.

Female. Head broader than the chest: feelers subclavate, longer than the chest; joints from the fifth to the tenth successively decreasing in length: abdomen oval, convex above, keeled beneath, as broad as the chest, but much shorter; metapodeon occupying more than half the back; octoon rather large; ennaton and following segments very short.

LAMPROTATUS Pycnos. Mas. *Cyaneo-viridis, abdomine cupreo vario, antennis nigris, tibiis fulvis apice tarsisque piceis, alis subgriseis*. Page 111.

Bluish green: head and chest convex, shagreened: head a little broader than the chest: eyes and eyelets red: feelers black, filiform, longer than the chest; first joint green, long, slender; second cup-shaped; third and fourth very small; fifth and following joints successively decreasing in length; club spindle-shaped, much longer than the tenth joint: chest oval: fore-chest short; its breadth above more than twice its length: sutures of the parapsides distinct; axillæ large, approaching near to each other; scutcheon subrhomboidal, with a very slight transverse suture near its tip: hind-chest blue, rather large, obconical, declining: petiole very short: abdomen smooth, flat, nearly linear, contracted at the base, varied with blue and copper-colour, shorter and rather narrower than the chest; metapodeon occupying about half of the back; octoon and ennaton of moderate size; decaton and following segments short: legs tawny; hips and thighs green, the latter and especially the fore-thighs tawny towards their tips; middle-shanks and hind-shanks with piceous tips, each having also a broad brown band near the base; feet piceous, hind-feet somewhat paler than the other feet: wings with a very slight gray tinge; ulna rather more than half the length of the humerus; radius much longer than the ulna; cubitus more than half the length of the ulna; wing-brand of moderate size. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

Var. *a*. Head and chest green: abdomen brassy brown: no bands across the hinder-shanks.

Female ? Greenish blue: head a little broader than the chest: feelers clavate, a little longer than the chest; joints from the fifth to the tenth successively decreasing in length; club nearly elliptical, much longer than the tenth joint: abdomen elliptical, smooth, purple, blue at the base, convex above, keeled beneath, much shorter but not narrower than the chest; metapodeon occupying one-half of the back; octoon and ennaton of moderate size; decaton and following segments very short.

LAMPROTATUS Coretas. Mas. *Cyaneo-viridis, abdominis disco purpureo-cupreo, antennis nigris, tibiis tarsisque piceis, protibiis fulvis piceo vittatis, alis subfuscis.* Page 111.

Dark green, with a slight bluish tint: head and chest convex, shagreened: head broader than the chest: eyes and eyelets dark red: feelers black, filiform, a little longer than the chest; first joint green; second cup-shaped; third and fourth very small; fifth and following joints to the tenth successively decreasing in length; club spindle-shaped, very much longer than the tenth joint: chest oval: fore-chest short; its breadth above about twice its length: sutures of the parapsides distinct; axillæ rather large, parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a distinct transverse suture near its tip: hind-chest rather large, obconical, somewhat rugulose, slightly declining: petiole very short: abdomen linear, smooth, depressed, purplish copper on the disk, shorter and very much narrower than the chest; metapodeon occupying most of the back; octoon and following segments short: legs piceous; hips and thighs green; knees and fore-shanks tawny, the latter having a piceous line along their whole length: wings rather narrow, with a very slight brownish tint; veins brown; ulna about half the length of the humerus; radius much longer than the ulna; cubitus full half the length of the ulna; wing-brand piceous, rather small. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

SEMIOTUS Tauriscus. Mas. *Viridis, abdominis disco purpureo, antennis nigris, pedibus piceis, femoribus viridibus, protibiis et protarsis fulvis, alis subgriseis.* Page 112.

Dark green: head and chest convex, finely shagreened: head not broader than the chest: eyes and eyelets dark red: feelers black, filiform, not longer than the chest; first joint long, slender, green; second cup-shaped; third and fourth very small; the fol-

lowing joints from the fifth to the tenth successively decreasing in length; club long-conical, much longer than the tenth joint: chest elliptical: fore-chest very short; its length above not more than one-eighth of its breadth; hind-border very concave: shield of the mid-chest rather long; sutures of the parapsides distinct; axillæ parted by nearly one-third of the breadth of the chest; scutcheon subrhomboidal: hind-chest rather broad, short-obconical, slightly declining: petiole very short: abdomen oval, smooth, depressed, dark purple on the disk, a little shorter but hardly narrower than the chest; metapodeon occupying more than half of the back; octoon about one-third of the length of the metapodeon; ennaton shorter; decaton and the following segments very short: legs tawny; hips and thighs green; four hinder-shanks and feet piceous: wings very slightly tinged with gray; veins piceous; ulna a little shorter than the humerus; radius shorter than the ulna; cubitus about half the length of the radius; wing-brand very small, forked. Length of the body 1 line; of the wings $1\frac{3}{4}$ line.

PTEROMALUS *Æson*. Mas. *Viridis, abdomine purpureo-æneo, antennis fuscis, pedibus flavis, femoribus viridibus, alis subfulvis.* Page 119.

Body nearly linear: head and chest green, convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers brown, filiform, as long as the chest; first joint yellow, long, linear; second cup-shaped; third and fourth very small; fifth and following joints oblong-quadrate, but successively decreasing in length; club conical, more than twice the length of the tenth joint: chest nearly linear: fore-chest very short: shield of the middle-chest rather long; sutures of the parapsides indistinct; axillæ separated by rather less than one-fourth of the breadth of the chest; scutcheon subrhomboidal, having a very slight trace of a transverse suture before its tip: hind-chest of moderate size, obconical, declining: petiole very short: abdomen æneous-green, flat, smooth, shining, shorter than the chest; the disk is purplish brown; metapodeon occupying nearly one-third of the back; octoon and following segments of moderate size: legs bright yellow; hips and thighs green; tips of the feet brown: wings with a slight tawny tinge, especially beneath the ulna and the radius; ulna hardly half the length of the humerus; radius a little longer than the ulna; cubitus shorter than the ulna; wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

PTEROMALUS Camma. Fem. *Viridis, abdomine æneo, basi viridi-fulvo, antennis nigris, pedibus flavis, alis limpidis.*
Page 119.

Body convex: head and chest dark green, finely shagreened: head broader than the chest: eyes and eyelets red: feelers black, very slightly clavate, much longer than the chest; first joint long, slender, linear, yellow; second cup-shaped, piceous; third and fourth very small; fifth and following joints oblong, successively decreasing in length; club spindle-shaped, piceous, hardly broader than the tenth joint but more than twice its length: chest elliptical: fore-chest very short: shield of the middle-chest of moderate size; sutures of the parapsides very indistinct; axillæ separated by one-fourth of the breadth of the chest; scutcheon nearly rhomboidal: hind-chest of moderate size, obconical, declining: petiole short: abdomen elliptical, bronze, smooth, shining, greenish tawny at the base, pointed at the tip, rather deeply keeled beneath, a little narrower and longer than the chest; metapodeon rather large; octoon and following segments shorter: oviduct yellow: legs yellow; hips green; tips of feet brown: wings colourless; veins pale yellow; ulna very much shorter than the humerus; radius longer than the ulna; cubitus shorter than the ulna; wing-brand very small. Length of the body $\frac{2}{3}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Phasis. Mas. *Cyaneo-viridis, abdominis disco purpureo-cupreo, antennis tibiisque piceis, femoribus viridibus, tarsis flavis, alis limpidis.* Page 119.

Head and chest bluish green, convex, finely shagreened: head a little broader than the chest; front blue: eyes and eyelets piceous: feelers filiform, piceous, longer than the chest; first joint long, slender, green; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth linear, of nearly equal size; club spindle-shaped, more than twice the length of the tenth joint: chest elliptical: fore-chest very short; its length above not more than one-tenth of its breadth; hind-border concave: shield of the mid-chest broad; sutures of the parapsides indistinct; axillæ parted by nearly one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest broad, declining, with a ridge along the middle and a rim on each side; petiole very short: abdomen green, purplish copper on the disk, flat, smooth, increasing in breadth from the base till near the tip, a little narrower and much shorter than the chest; metapodeon occupying nearly one-third of the back; octoon and following segments of moderate length: hips and thighs green;

shanks piceous ; knees, feet, and tips of shanks yellow ; tips of four hinder feet brown : wings colourless ; veins pale tawny ; ulna about half the length of the humerus ; radius a little shorter than the ulna ; cubitus a little shorter than the radius ; wing-brand very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Cyamon. Mas. *Viridis, abdomine purpureo, antennis pedibusque fulvis, femoribus piceo vittatis, alis limpidis.* Page 119.

Head and chest bright green, convex, finely shagreened : head a little broader than the chest : feelers dark tawny, subclavate, as long as the chest ; first joint tawny, long, linear ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club elliptical, broader than the tenth joint and more than twice its length : chest elliptical, rather long, with a slight brassy tinge : fore-chest very short ; its length above not more than one-tenth of its breadth ; hind-border concave : shield of mid-chest rather long ; sutures of the parapsides very indistinct ; axillæ parted by about one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest short, declining, with a rim on each side : petiole very short : abdomen linear, flat, smooth, dark purple, a little narrower and much shorter than the chest ; metapodeon occupying more than one-third of the back ; octoon rather long ; the following segments short : legs pale tawny ; hips green ; thighs with a piceous line above, which line in the hind-pair extends over the whole surface of the thighs ; middle-feet and hind-feet yellow with brown tips : wings colourless ; veins pale yellow ; ulna about half the length of the humerus ; radius as long as the ulna ; cubitus much shorter than the ulna ; wing-brand very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Hyloe. Fem. *Viridis, abdomine æneo-purpureo basi viridi micante, antennis nigris, pedibus luteis, femoribus viridibus, alis subfulvis.* Page 119.

Head and chest convex, finely shagreened : head dark green, a little broader than the chest : eyes and eyelets red : feelers black, clavate, rather stout, not longer than the chest ; first joint long, tawny, black at the tip ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club long-conical, not broader than the tenth joint but more than twice its length : chest nearly spindle-

shaped : fore-chest very short ; its length above less than one-tenth of its breadth ; hind-border concave : sutures of the parapsides indistinct ; scutcheon subrhomboidal, with a suture along each side : hind-chest of moderate size, obconical, declining : petiole very short : abdomen oval, smooth, coppery-purple, bright green at the base, depressed above, slightly keeled beneath, shorter but not broader than the chest ; metapodeon occupying about one-fourth of its length ; octoon and following segments of moderate and nearly equal size : legs bright luteous ; hips and thighs green ; four hinder feet yellow with piceous tips : wings slightly tawny ; veins tawny ; ulna more than half the length of the humerus ; radius longer than the ulna ; cubitus much shorter than the ulna ; wing-brand very small. Length of the body $1\frac{1}{2}$ line ; of the wings $2\frac{3}{4}$ lines.

PTEROMALUS Acco. Fem. *Æneo-viridis, abdomine purpureo æneo, basi viridi, antennis nigris, femoribus æneis, alii fulvis.* Page 120.

Allied to *P. deplanatus*. Body dull brassy green : head and chest convex, finely shagreened : head hardly broader than the chest : eyes and eyelets red : feelers black, clavate, rather shorter than the chest ; first joint tawny, very long ; second long-cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth short, closely joined together and successively decreasing in length ; club conical, more than twice the length of the tenth joint : chest nearly elliptical : fore-chest very short ; its length above less than one-tenth of its breadth ; hind-border rather concave : shield of the mid-chest rather short and broad ; sutures of the parapsides very indistinct ; axillæ parted by about one-fourth of the breadth of the chest ; scutcheon subrhomboidal, with a suture along each side : hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen nearly round, smooth, purplish bronze, green at the base, depressed above, hardly keeled beneath, much shorter than the chest, its length scarcely exceeding its breadth ; metapodeon occupying hardly one-third of the back ; octoon and two following segments of moderate length ; the rest very short : legs tawny ; hips and thighs brassy : wings tawny, rather narrow ; veins dull tawny ; ulna hardly half the length of the humerus ; radius a little shorter than the ulna ; cubitus shorter than the radius ; wing-brand very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Tychon. Fem. *Niger, abdomine aeneo macula basali fulva, antennis nigris, pedibus fulvis, alis limpidis.*
Page 120.

Body very broad: head and chest black, convex, shagreened: head large, broader than the chest: eyes and eyelets dark red: feelers black, subclavate, much longer than the chest; first joint long, slender, tawny; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club spindle-shaped, a little broader than the tenth joint and more than twice its length: chest short-elliptical, robust: fore-chest extremely short; its length above less than one-tenth of its breadth; hind-border slightly concave: shield of the mid-chest broad and short; sutures of the parapsides very indistinct; axillæ parted by nearly one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a slight transverse suture near its tip: hind-chest large, declining, slightly obconical, with a ridge along the middle: petiole very short: abdomen short-elliptical, smooth, flat above, keeled beneath, brassy with an indistinct tawny spot near the base which is green, much shorter but hardly narrower than the chest; its length slightly exceeding its breadth; metapodeon occupying about one-third of the back; octoon full half the length of the metapodeon; ennaton and all the following segments short but distinct: legs tawny; hips brassy: wings colourless; veins tawny; ulna rather less than half the length of the humerus; radius a little longer than the ulna; cubitus much shorter than the ulna; wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{4}$ lines.

PTEROMALUS Cerycus. Fem. *Viridis, abdomine cyaneo purpureoque vario, antennis nigris, pedibus fulvo-flavis, alis limpidis.* Page 120.

Body bright green, rather narrow: head and chest convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers black, clavate, slender, nearly as long as the chest; first joint long, slender, yellow, slightly bent, black towards the tip; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club conical, more than twice the length of the tenth joint: chest nearly elliptical: fore-chest very short; its length above less than one-tenth of its breadth; hind-border concave: shield of the mid-chest rather flat; sutures of the parapsides very indistinct; axillæ parted by nearly one-third of the breadth of the chest; scutcheon

subrhomboidal : hind-chest rather large, obconical, declining, with a ridge along the middle and a short suture on each side : petiole very short : abdomen nearly oval, smooth, bluish green, dark coppery-purple on the disk, depressed above, hardly keeled beneath, shorter and a very little broader than the chest ; metapodeon occupying less than one-fourth of the back ; octoon and all the following segments of moderate and nearly equal size : legs bright tawny ; hips green ; shanks and feet of four hinder-legs bright yellow ; tips of the feet brown : wings colourless ; veins pale yellow ; ulna much more than half the length of the humerus ; radius a little shorter than the ulna ; cubitus much shorter than the radius ; wing-brand small, brown. Length of the body $1\frac{1}{2}$ line ; of the wings $2\frac{1}{2}$ lines.

PTEROMALUS Arestor. Fem. *Viridis, capite cyaneo-nigro, abdomine aeneo-viridi, antennis nigris, tibiis tarsisque flavis, alis limpidis.* Page 120.

Body rather narrow : head and chest convex, finely shagreened : head bluish black, broader than the chest : eyes and eyelets dark red : feelers black, slightly clavate, nearly as long as the chest ; first joint long, slender ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club spindle-shaped, more than twice the length of the tenth joint : chest bright green, nearly spindle-shaped : fore-chest very short ; its length not more than one-tenth of its breadth ; hind-border concave : shield of the mid-chest rather long ; sutures of the parapsides very indistinct ; axillæ parted by near one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest large, obconical, slightly declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen elliptical, brassy green, smooth, depressed above, hardly keeled beneath, a little broader and much shorter than the chest ; metapodeon occupying nearly one-fourth of the back ; octoon and the following segments of moderate size : legs bright yellow ; hips and thighs green, tips of the latter yellow ; tips of the feet brown : wings colourless ; veins pale yellow ; ulna rather more than half the length of the humerus ; radius shorter than the ulna ; cubitus shorter than the radius ; wing-brand very small. Length of the body 1 line ; of the wings 2 lines.

PTEROMALUS Eulimene. Fem. *Viridis, abdomine cupreo-purpureo, basi viridi micante, antennis pedibusque fulvis, femoribus viridibus, alis limpidis.* Page 120.

Body short, robust : head and chest dark green, convex, finely

shagreened: head as broad as the chest: eyes and eyelets red: feelers very dark tawny, subclavate, longer than the chest; first joint tawny, long and slender; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club elliptical, broader than the tenth joint and rather more than twice its length: chest broad, very short-elliptical: fore-chest rather short; its length above about one-sixth of its breadth: shield large; sutures of the parapsides indistinct; axillæ parted by rather less than one-third of its breadth; scutcheon subrhomboidal, with a suture along each side: hind-chest transverse, very short, bright green, almost smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen oval, smooth, shining, coppery-purple, bright green at the base, depressed above, rather deeply keeled beneath, a little narrower but hardly longer than the chest; metapodeon occupying rather more than one-fourth of the back; octoon and the three following segments rather short, nearly equal; paratelum longer; telum very small: legs pale tawny; hips and thighs green; four hinder-feet pale yellow with brown tips: wings colourless; veins tawny; ulna less than half the length of the humerus; radius a little longer than the ulna; cubitus shorter than the ulna; wing-brand brown, small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Dipœnos. Fem. *Æneo-viridis, abdomine æneo, antennis nigris, pedibus fulvis, femoribus piceis, alis subfulvis.* Page 120.

Head and chest brassy green, convex, finely shagreened: head very little broader than the chest: eyes and eyelets piceous: feelers black, clavate, as long as the chest; first joint dark tawny, long and slender; second cup-shaped, piceous; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club conical, rather broader than the tenth joint and more than twice its length: chest nearly elliptical: fore-chest very short; its length above less than one-tenth of its breadth: shield of the mid-chest broad and short; sutures of the parapsides very indistinct; axillæ parted by less than one-third of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest rather short, obconical, declining: petiole very short: abdomen oval, smooth, bronze-colour, bright coppery-green at the base, depressed above, rather deeply keeled beneath, a little longer but hardly broader than the chest; metapodeon occupying about one-fourth of the back; octoon shorter than the metapodeon; ennaton still shorter; decaton as long as the octoon; protelum

shorter; paratelum and telum very short: legs pale tawny; hips brassy; thighs and tips of four hinder-feet piceous: wings with a slight tawny tinge; veins tawny; ulna full half the length of the humerus; radius a little shorter than the ulna; cubitus much shorter than the radius; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Glautias. Fem. *Viridis, abdomine purpureo basi aureo-viridi, antennis piceis, pedibus flavis, femoribus viridibus, tibiis fusco cinctis, alis limpidis.* Page 120.

Bright green: head and chest convex, finely shagreened: head a little broader than the chest: eyes and eyelets dark red: feelers dark piceous, subclavate, rather slender, hardly longer than the chest; first joint green, long, slender; second cup-shaped, third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club nearly elliptical, rather broader than the tenth joint and more than twice its length; chest short-elliptical: fore-chest short; its length above about one-eighth of its breadth; hinder-border slightly concave; shield of the mid-chest broad; sutures of the parapsides very indistinct; axillæ parted by one-third of the breadth of the chest; scutcheon sub-rhomboidal, with a suture along each side: hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, smooth, purple on the disk, golden green at the base, depressed above, keeled beneath, a little narrower and much longer than the chest; metapodeon occupying hardly one-fifth of the back; octoon and enaton of nearly equal size; decaton, protelum and paratelum longer; telum short: legs yellow; hips and thighs green; a brown band round the base of each shank; four hinder-feet pale yellow with piceous tips: wings colourless; veins pale yellow; ulna less than half the length of the humerus; radius much longer than the ulna; cubitus shorter than the ulna; wing-brand small, pale brown. Length of the body $1\frac{1}{2}$ line; of the wings $2\frac{1}{2}$ lines.

PTEROMALUS Suia. Mas. *Cyaneo- aut aureo-viridis, abdominis disco purpureo, antennis pedibusque fulvis, femoribus viridibus, alis limpidis.* Page 120.

Head and chest bright bluish green, convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers dark tawny, clavate, rather slender, a little longer than the chest; first joint long, slender, tawny, black at the tip; second black, cup-

shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club spindle-shaped, broader than the tenth joint and more than twice its length: chest oval: fore-chest very short; its length less than one-tenth of its breadth: shield of the mid-chest rather long; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short; abdomen nearly linear, depressed, smooth, green, purple on the disk, a little shorter and narrower than the chest; metapodeon occupying about one-fourth of the back; octoon and following segments short: legs tawny; hips and thighs excepting the tips of the latter green; four hinder-knees and feet pale tawny; tips of the latter brown: wings colourless; veins pale tawny; ulna less than half the length of the humerus; radius much longer than the ulna; cubitus shorter than the radius; wing-brand small, pale brown. Length of the body 1 line; of the wings 2 lines.

Var. *a*. Golden green; disk of the abdomen purple: wing-brand pale tawny.

PTEROMALUS Amnisos. Fem. Æneus, abdomine purpureo basi viridi, antennis nigris apice piceis, pedibus flavis, femoribus viridibus, tibis picco cinctis, alis limpidis. Page 120.

Body brass colour: head and chest convex, finely shagreened: head as broad as the chest: eyes and eyelets dark red: feelers slender, subclavate, black, piceous at the tips, as long as the chest; first joint long, slender, tawny at the base; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club conical at the tip, a little broader than the tenth joint and more than twice its length: chest oval: fore-chest short; its length rather more than one-tenth of its breadth: sutures of the parapsides very indistinct; axillæ parted by rather more than one-fourth of the chest; scutcheon subrhomboidal, with a suture along each side and an indistinct transverse channel near the tip: hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen long-oval, smooth, purple on the disk, bright green at the base, depressed above, deeply keeled beneath, as long and nearly as broad as the chest; metapodeon occupying less than one-fourth of the back; hind-border convex; octoon and following segments of nearly equal size: legs yellow; hips and thighs green; a broad piceous band across each of the four hinder-thighs;

fore-shanks and fore-feet tawny ; four hinder-feet with piceous tips : wings colourless ; veins tawny ; ulna hardly half the length of the humerus ; radius much longer than the ulna ; cubitus slightly curved, shorter than the ulna. Length of the body $1\frac{1}{2}$ line ; of the wings $2\frac{3}{4}$ lines.

PTEROMALUS Zagreus. Fem. Viridis, abdomine cyaneo purpureoque vario, antennis nigris, pedibus flavis, femoribus viridibus, alis limpidis. Page 120.

Body narrow : head and chest convex, green, finely shagreened : head broader than the chest : eyes and eyelets dark red : feelers black, slender, clavate, not longer than the chest ; first joint long, slender, green, yellow at the base ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club nearly conical, much broader than the tenth joint, and more than twice its length : chest long, elliptical : fore-chest very short ; its length full one-tenth of its breadth : shield of the mid-chest rather long ; sutures of the parapsides indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal, with a suture along each side : hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen spindle-shaped, smooth, dark green, tinged with blue and purple, depressed above, slightly keeled beneath, longer and a little broader than the chest ; metapodeon occupying about one-fifth of the back ; octoon and ennaton of moderate and nearly equal size ; decaton, protelum, and paratelum longer ; telum very small : legs bright yellow ; hips and thighs green ; tips of feet brown ; fore-shanks and fore-feet tawny : wings colourless ; veins pale yellow ; ulna not half the length of the humerus ; radius a little shorter than the ulna ; cubitus a little shorter than the radius ; wing-brand pale brown, very small. Length of the body 1 line ; of the wings 2 lines.

PTEROMALUS Opheltes. Mas. Viridis, abdomine æneo, antennis nigris, pedibus flavis, femoribus viridibus, alis limpidis. Page 120.

Head and chest dark green, convex, finely shagreened : head broader than the chest : eyes and eyelets piceous : feelers black, subclavate, much longer than the chest ; first joint green, long, slender ; second cup-shaped ; third and fourth very small ; the fol-

lowing joints to the tenth successively decreasing in length; club spindle-shaped, rather broader than the tenth joint and nearly twice its length: chest short-elliptical: fore-chest very short; its length not one-tenth of its breadth: shield of the mid-chest rather flat; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest obconical, declining, nearly smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen nearly linear, depressed, smooth, bronze colour, tinged with green at the base, a little longer and very much narrower than the chest; metapodeon occupying more than one-third of the back: legs pale yellow; hips and thighs dark green; fore-shanks and fore-feet tawny; tips of four hinder-feet brown; wings colourless; veins tawny; ulna rather thick, full half the length of the humerus; radius nearly as long as the ulna; cubitus much shorter than the radius; wing-brand pale brown, very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Mese. Mas. *Aureus*, abdomine purpureo, basi viridi flavoque maculato, antennis fulvis, pedibus flavis, alis limpidis. Page 120.

Body golden colour, rather narrow: head and chest convex, finely shagreened: head broader than the chest: eyes and eyelets dark red: feelers tawny, subclavate, a little shorter than the chest; first joint long, slender, bright pale yellow; second piceous, cup-shaped; third and fourth very small; the following joints to the tenth of equal breadth but successively decreasing in length; club spindle-shaped, broader than the tenth joint and more than twice its length: chest nearly spindle-shaped: fore-chest very short; its length less than one-tenth of its breadth: parapsides very indistinct; axillæ parted by less than one-fourth of the breadth of the chest; scutcheon rather long, subrhomboidal: hind-chest well developed, obconical, declining, nearly smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen nearly linear, depressed, smooth, purple, a little narrower and much shorter than the chest, having a large yellow spot near the base which is bright green; metapodeon occupying nearly one-third of the back; octoon and following segments of nearly equal size: legs very bright pale yellow; hips gold colour: wings colourless; veins very pale tawny; ulna full half the length of the humerus; radius as long as the ulna; cubitus much shorter than the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Alopius. Mas. *Viridis, abdominis disco cupreo purpureo, antennis fulvis, pedibus flavis, alis limpidis*. Page 120.

Very bright green : head and chest convex, finely shagreened : head much broader than the chest : eyes and eyelets piceous : feelers tawny, slender, subclavate, a little shorter than the chest ; first joint long, slender, pale yellow, luteous at the tip ; second long-cup-shaped ; third and fourth very small ; the following joints to the tenth successively decreasing in length ; club spindle-shaped, a little broader than the tenth joint and more than twice its length : chest nearly elliptical : fore-chest very short ; its length not more than one-tenth of its breadth : sutures of the parapsides indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal, with a suture along each side : hind-chest golden green, short-obconical, declining, almost smooth, with a ridge along the middle and a rim on each side : petiole very short : abdomen depressed, smooth, increasing in breadth from the base till near the tip, hardly shorter or narrower than the chest ; disk coppery purple ; metapodeon occupying nearly one-third of the back ; octoon and following segments of nearly equal size : legs bright yellow ; hips green ; fore-shanks and fore-feet bright tawny ; tips of four hinder-feet brown : wings colourless ; veins pale tawny ; ulna about half the length of the humerus ; radius a little longer than the ulna ; cubitus shorter than the ulna ; wing-brand small, pale brown. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Phylacis. Mas. *Niger, tarsi flavis, alis limpidis*. Page 120.

Black : head and chest convex, finely shagreened : head much broader than the chest : eyes and eyelets dark red : feelers black, slender, filiform, longer than the chest ; first joint long, slender ; second cup-shaped ; third and fourth very small ; the following joints to the tenth successively decreasing in length ; club long-spindle-shaped, broader than the tenth joint and more than twice its length : chest rather long, nearly elliptical : fore-chest rather short ; its length about one-sixth of its breadth : shield rather long ; sutures of the parapsides indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal, with a slight impression at the base : hind-chest well developed, long, obconical, declining, with a slight ridge along the middle : petiole very short : abdomen elliptical, flat, smooth, very much shorter and narrower than the chest ; metapodeon occupying one-half of the back ; octoon

and following segments short: legs black; knees and fore-feet tawny; four hinder-feet yellow with brown tips: wings colourless; veins tawny; ulna more than half the length of the humerus; radius not longer than the ulna; cubitus much shorter than the radius; wing-brand extremely small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS CÆNO. Mas. *Aureo-viridis, abdominis disco purpureo æneo, antennis nigris, pedibus fulvis, femoribus viridibus, alis limpidis.* Page 120.

Golden green: head and chest convex, finely shagreened: head hardly broader than the chest: eyes and eyelets dark red: feelers black, filiform, a little longer than the chest; first joint long, slender, yellow at the base; second cup-shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest nearly oval: fore-chest very short; its length hardly more than one-tenth of its breadth: sutures of the parapsides rather indistinct, especially towards the hind-border of the shield; axillæ as usual with a suture along the middle, parted by full one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side, and a slight transverse channel near the tip: hind-chest rather short, slightly obconical, declining, almost smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen smooth, depressed, slightly increasing in breadth from the base till near the tip, very little shorter and narrower than the chest; disk purplish bronze; segments of nearly equal size: sexual parts brown: legs tawny; hips and thighs green; four hinder-knees and feet pale tawny; tips of the latter brown: wings colourless; veins tawny; ulna not less than half the length of the humerus; radius longer than the ulna; cubitus shorter than the ulna; wing-brand small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{4}$ lines.

Var. β . Abdomen bluish green.

PTEROMALUS CORION. Fem. *Cyaneus, abdomine purpureo-æneo, basi viridi micante, antennis pedibusque anticis fulvis, femoribus piceis, tibiis fuscis, tarsis flavis, alis limpidis.* Page 120.

Head and chest blue, convex, finely shagreened: head a little broader than the chest: eyes and eyelets dark red: feelers tawny, slender, subclavate, a little longer than the chest; first joint pale

tawny, long and slender; second cup-shaped; third and fourth very small; the following joints from the fifth to the ninth successively decreasing in length: club spindle-shaped, rather broader than the tenth joint and more than twice its length: chest broad, short-elliptical: fore-chest very short; its length less than one-tenth of its breadth: shield of the mid-chest large; sutures of the parapsides very indistinct, almost obsolete; axillæ parted by less than one-third of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest transverse, very short, declining, bright green, almost smooth: petiole very short: abdomen oval, smooth, purplish bronze, bright green at the base, depressed above, slightly keeled beneath, much longer but not broader than the chest; segments of moderate and nearly equal size: legs tawny; hips blue; thighs and tips of feet piceous; four hinder-legs with brown shanks and pale yellow feet: wings colourless; veins tawny; ulna less than half the length of the humerus; radius none; cubitus much shorter than the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS *Hermachus*. Mas. *Cupreus, abdomine basi apice-que viridi, antennis nigris, pedibus flavis, femoribus viridibus, alis limpidis.* Page 120.

Copper colour: head and chest convex, finely shagreened, dark green beneath: head much broader than the chest: eyes and eyelets dark red: feelers subclavate, black, rather longer than the chest; first joint green, long, slender; second cup-shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club spindle-shaped, a little broader than the tenth joint and more than twice its length: chest long-oval; fore-chest very short; its length not more than one-tenth of its breadth: sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest green, nearly smooth, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen elliptical, depressed, smooth, green towards the tip, bright green at the base, shorter and narrower than the chest; metapodeon occupying about one-third of the back: legs bright yellow; hips and thighs green; tips of four hinder-feet pale brown: wings colourless; veins tawny; ulna about half the length of the humerus; radius very short; cubitus full twice the length of the radius, and not much shorter than the ulna; wing-brand very small. Length of the body $\frac{2}{3}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Temesa. Mas. *Viridis, abdomine flavo maculato, antennis luteis, pedibus flavis, alis limpidis.* Page 120.

Bright green : head and chest convex, finely shagreened : head a little broader than the chest : eyes and eyelets red : feelers sub-clavate, luteous, longer than the chest ; first joint pale yellow, long, slender ; second cup-shaped ; third and fourth very small ; the following joints from the tenth successively decreasing in length ; club spindle-shaped, a little broader than the tenth joint and nearly thrice its length : chest nearly spindle-shaped : fore-chest very short ; its length above not one-tenth of its breadth : shield of the mid-chest rather flat ; sutures of the parapsides very indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon copper colour, rhomboidal : hind-chest well developed, obconical, declining, nearly smooth, with a ridge along the middle and a rim on each side : petiole very short : abdomen linear, depressed, smooth, having an indistinct yellow spot near the base, narrower and a little longer than the chest ; metapodeon occupying about one-third of the back ; octoon and following segments nearly equal in size : legs bright pale yellow ; hips green : wings colourless ; veins pale yellow ; ulna hardly half the length of the humerus ; radius longer than the ulna ; cubitus much shorter than the ulna ; wing-brand very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Drepanon. Mas. *Cyaneo-viridis, abdomine basi fulvo, antennis femoribusque piceis, pedibus fulvis, alis sublimpidis.* Page 120.

Dark bluish green : head and chest convex, finely shagreened : head hardly broader than the chest : eyes and eyelets piceous : feelers piceous, filiform, a little longer than the chest ; first joint tawny, long, slender ; second cup-shaped ; third and fourth very small ; the following joints to the tenth successively decreasing in length ; club spindle-shaped, more than twice the length of the tenth joint : chest elliptical : fore-chest very short ; its length above not more than one-tenth of its breadth : shield of the mid-chest rather flat ; sutures of the parapsides very indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest of moderate size, obconical, declining, with a ridge along the middle, and a rim on each side : petiole very short : abdomen slightly increasing in breadth from the base till near the tip, depressed, smooth, tawny towards the base, nearly as long and as broad as the chest : sexual parts pale brown : legs tawny ; hips

black; thighs piceous; four hinder-knees and feet yellow: wings nearly colourless; veins tawny; ulna rather thick, much more than half the length of the humerus; radius a little longer than the ulna; cubitus hardly half the length of the ulna; wing-brand small. Length of the body $\frac{1}{2}$ line; of the wings 1 line.

PTEROMALUS Belesis. Mas. Viridis, abdomine æneo basi viridi, antennis fulvis, pedibus luteis, alis limpidis. Page 121.

Head and chest bright green: head a little broader than the chest: eyes and eyelets dark red: feelers tawny, subclavate, shorter than the chest; first joint luteous, long, slender; second cup-shaped; third and fourth very small; the following joints to the tenth short and successively decreasing in length; club conical, a little broader than the tenth joint, and more than twice its length: chest nearly oval: fore-chest very short; its length not one-tenth of its breadth: shield of the mid-chest short; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest: hind-chest transverse, very short: petiole very short: abdomen elliptical, depressed, smooth, bronze colour, bright green at the base, shorter and a little narrower than the chest; metapodeon occupying less than one-third of the back; octoon and following segments of nearly equal size: legs luteous; hips green; four hinder-feet yellow with brown tips: wings colourless; veins dull yellow; ulna much less than half the length of the humerus; radius a little longer than the ulna; cubitus as long as the ulna; wing-brand very small. Length of the body $\frac{2}{3}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Lebadeia. Fem. Viridi-cupreus, antennis nigris, pedibus fulvis, alis limpidis. Page 121.

Copper colour, tinged with green: head and chest convex, finely shagreened: head much broader than the chest: eyes and eyelets dark red: feelers black, subclavate, not longer than the chest; first joint long, slender, tawny at the base; second cup-shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club spindle-shaped, a little broader than the tenth joint and more than thrice its length: chest short-oval: fore-chest very short; its length not one-tenth of its breadth; shield of the mid-chest broad; sutures of the parapsides very indistinct; axillæ parted by rather more than one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest obconical, declining, nearly smooth,

with a ridge along the middle and a rim on each side : petiole very short : abdomen spindle-shaped, smooth, depressed above, hardly keeled beneath, much narrower and a little longer than the chest ; metapodeon occupying nearly half the back ; octoon and following segments short and of nearly equal length : legs tawny ; hips green ; four hinder-feet yellow with brown tips : wings colourless ; veins pale tawny ; ulna nearly half the length of the humerus ; radius much longer than the ulna ; cubitus a little shorter than the ulna ; wing-brand pale brown, very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Bercynthos. Mas. Cyaneo-viridis, abdominis disco purpureo-cupreo, antennis fulvis, pedibus luteis, alis subfulvis. Page 121.

Head and chest convex, finely shagreened : head bright blue, a little broader than the chest : eyes and eyelets red : feelers tawny, subclavate, not longer than the chest ; first joint long, slender, luteous ; second cup-shaped ; third and fourth very small ; the following joints to the tenth successively decreasing in length ; club at the tip a little broader than the tenth joint and more than twice its length : chest oval, bright bluish green : fore-chest very short ; its length above not one-tenth of its breadth : shield of the mid-chest broad ; sutures of the parapsides very indistinct ; axillæ parted by rather more than one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest short, declining : petiole very short : abdomen linear, depressed, smooth, green, hardly narrower or shorter than the chest ; disk purplish copper ; metapodeon occupying more than half the length of the back : legs luteous ; hips green ; four hinder-feet yellow with tawny tips : wings with a slight tawny tinge ; veins tawny ; ulna less than half the length of the humerus ; radius much longer than the ulna ; cubitus as long as the ulna ; wing-brand very small. Length of the body 1 line ; of the wings 2 lines.

Var. β . Head and chest green.

PTEROMALUS Lebene. Fem. Viridis, abdominis disco æneo, antennis fulvis, pedibus luteis, femoribus viridibus, alis subfulvis. Page 121.

Bright green : head and chest convex, finely shagreened : head hardly broader than the chest : eyes and eyelets red : feelers tawny, subclavate, not longer than the chest ; first joint luteous, long, slender ; second cup-shaped, piceous ; third and fourth very small ; the

following joints to the tenth successively decreasing in length; club spindle-shaped, a little broader than the tenth joint and more than twice its length: chest nearly elliptical: fore-chest very short; its length less than one-tenth of its breadth: shield of the mid-chest short; sutures of the parapsides very indistinct; axillæ parted by nearly one-third of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest transverse, short, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen long-oval, smooth, depressed above, rather deeply keeled beneath, longer but not narrower than the chest; disk bronze colour; metapodeon occupying hardly one-fourth of the chest; octoon and following segments of equal size: legs luteous; hips and thighs, except their tips, green; tips of four hinder-feet brown: wings with a slight tawny tinge; veins yellow; ulna rather less than half the length of the humerus; radius a little shorter than the ulna; cubitus shorter than the radius; wing-brand very small. Length of the body 1 line; of the wings 2 lines.

PTEROMALUS Cercaphrus. Mas. *Æneo-viridis, abdomine viridi-æneo, antennis pedibusque fulvis, femoribus viridibus, alis limpidis.* Page 121.

Body short, broad: head and chest coppery-green, convex, finely shagreened: head large, broader than the chest: eyes and eyelets red: feelers tawny, clavate, a little shorter than the chest; first joint long, slender, green; second black, cup-shaped; third and fourth very small; the following joints to the tenth short and successively decreasing in length; club conical, a little broader than the tenth joint and more than twice its length: chest short-oval: fore-chest very short; its length above not more than one-tenth of its breadth: shield of the mid-chest short; sutures of the parapsides very indistinct; axillæ parted by one-fifth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side and a slight transverse channel near the tip: hind-chest transverse, short, declining, nearly smooth, with a slight ridge along the middle and a rim on each side: petiole very short: abdomen conical, depressed, smooth, much narrower than the chest and hardly more than half its length, bright copper-colour tinged with green; metapodeon occupying the greater part of the back: legs tawny; hips and thighs green; tips of shanks, knees and feet of four hinder-legs yellow; tips of feet brown: wings colourless; veins yellow; ulna about half the length of the humerus; radius nearly as long as the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body $\frac{1}{2}$ line; of the wings 1 line.

PTEROMALUS Lampe. Fem. *Viridis, abdomine aeneo-viridi disco purpureo-cupreo, antennis nigris, pedibus fulvis, alis limpidis.* Page 121.

Head and chest brassy, convex, finely shagreened: head hardly broader than the chest: eyes and eyelets dark red: feelers black, clavate, a little longer than the chest; first joint long, slender, tawny; second cup-shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club long-conical, a little broader than the tenth joint and more than twice its length: chest elliptical: fore-chest very short; its length above not one-tenth of its breadth: shield of the mid-chest rather short and broad; sutures of the parapsides very indistinct; axillæ parted by full one-third of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest of moderate size, obconical, declining, almost smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen long-elliptical, smooth, brassy green, bright green at the base, depressed above, hardly keeled beneath, much longer but not narrower than the chest; disk purplish bronze; metapodeon occupying less than one-fourth of the back; octoon and four following segments of nearly equal size; telum very short: legs tawny; hips brass colour; four hinder-feet yellow with brown tips: wings colourless; veins tawny; ulna about half the length of the humerus; radius longer than the ulna; cubitus much shorter than the ulna; wing-brand pale brown, very small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{4}$ lines.

PTEROMALUS Anchinoe. Mas. *Cupreus, abdomine basi viridi micante, antennis nigris, pedibus flavis, femoribus cupreis, alis limpidis.* Page 121.

Head and chest dark copper colour, convex, finely shagreened: head a little broader than the chest: eyes and eyelets dark red: feelers black, subclavate, a little longer than the chest; first joint very slender; second cup-shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club spindle-shaped, a little broader than the tenth joint and nearly thrice its length: chest elliptical: fore-chest very short; its length not more than one-tenth of its breadth: shield of the mid-chest rather flat above; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest smooth, declining, rather short: petiole very short: abdomen nearly linear, depressed, smooth, bright green

at the base, shorter and narrower than the chest; disk copper colour; metapodeon occupying less than one-third of the back; octoon, enaton, and decaton rather long; the following segments very short: legs yellow; hips and thighs dark copper colour; four hinder-feet with brown tips; fore-shanks and fore-feet pale tawny; wings colourless; veins tawny; ulna rather thick, more than half the length of the humerus; radius shorter than the ulna; cubitus shorter than the radius; wing-brand small, brown. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Gallonius. Mas. *Cyaneus, abdomine cupreo basi cyaneo-viridi, antennis nigris, pedibus fulvis, femoribus piceis, alis limpidis.* Page 121.

Head and chest blue, convex, very finely shagreened, so as to appear almost smooth: head much broader than the chest: eyes and eyelets piceous; feelers black, subclavate, very slender, a little longer than the chest; first joint long, slender, cup-shaped; third and fourth very small; the following joints to the tenth long, linear, but successively decreasing in length; club spindle-shaped, broader than the tenth joint and more than twice its length: chest nearly elliptical: fore-chest very short; its length above not more than one-tenth of its breadth: shield of the mid-chest long, rather flat; sutures of the parapsides indistinct; axillæ parted by less than one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest rather short, obconical, declining: petiole very short: abdomen nearly linear, depressed, smooth, copper colour, bluish-green at the base, a little shorter and narrower than the chest; metapodeon occupying more than one-third of the back; octoon and following segments of nearly equal size: legs tawny; hips blue; thighs piceous; four hinder-feet yellow with brown tips: wings colourless, rather small; veins tawny; ulna about half the length of the humerus; radius a little shorter than the ulna; cubitus nearly as long as the radius; wing-brand small. Length of the body 1 line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Eleuthera. Fem. *Æneo-viridis, abdomine purpureo-cupreo basi cyaneo-viridi, antennis nigris, pedibus rufis, alis subfulvis.* Page 121.

Head and chest convex, shagreened: head green, a little broader than the chest: eyes and eyelets dark red: feelers black, filiform, longer than the chest; first joint long, slender, tawny, piceous at the tip; second cup-shaped; third and fourth very small;

the following joints to the tenth successively decreasing in length; club long-conical, more than twice the length of the tenth joint; chest oval, coppery-green: fore-chest short; its length above less than one-tenth of its breadth; it is produced slightly in front where it is narrower, declining, and wrinkled: shield of the mid-chest broad; sutures of the parapsides indistinct; axillæ parted by rather less than one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side, and a well defined transverse channel near the tip: hind-chest rather short, obconical, declining, nearly smooth, with its fore-shield distinct, and having a forked ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, finely shagreened, purplish copper, bluish green at the base, depressed above, keeled beneath, hairy towards the tip, a little longer than the chest; metapodeon occupying less than one-sixth of the back; the following segments from the octoon to the paratelum successively increasing in length: legs red; hips green; feet yellow with tawny tips; sheaths of the oviduct black, hairy, appearing just beyond the tip of the abdomen: wings with a slight tawny tinge; each fore-wing with a large brown spot beneath the brand; veins piceous; ulna much less than half the length of the humerus; radius very much longer than the ulna; cubitus much shorter than the ulna; wing-brand small. Length of the body 2 lines; of the wings $3\frac{1}{4}$ lines.

PTEROMALUS Lyttus. Fem. *Æneo-viridis, abdomine cyaneo-viridi, disco purpureo, antennis nigris, pedibus fulvis, alis limpidis.* Page 121.

Head and chest brassy green, convex, finely shagreened: head a little broader than the chest: eyes and eyelets piceous: feelers black, clavate, rather hairy, much shorter than the chest; first joint long, slender, green; second cup-shaped; third and fourth very small; the following joints to the tenth successively decreasing in length; club long-conical, rather broader than the tenth joint and more than twice its length: chest long-oval: fore-chest very short; its length hardly more than one-tenth of its breadth: shield of the mid-chest broad; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest of moderate size, obconical, declining, nearly smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen long-elliptical, smooth, bluish-green, purple on the disk, bright green at the base, depressed above, slightly keeled beneath, longer and a little broader than the chest; metapodeon occupying rather less than one-fourth of the back; octoon and all the following segments

of nearly equal length: legs tawny; hips green; four hinder-feet yellow, with piceous tips: wings colourless; veins tawny; ulna a little more than half the length of the humerus; radius hardly longer than the ulna; cubitus much shorter than the ulna; wing-brand brown, small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{4}$ lines.

PTEROMALUS Bienna. Mas. *Cyaneo-viridis, abdomine purpureo-cupreo, antennis picetis, femoribus viridibus, alis limpidis.*
Page 125.

Head and chest convex, finely shagreened: head greenish blue, a little broader than the chest: eyes and eyelets red: feelers piceous, nearly filiform, not longer than the chest; first joint long, slender, tawny towards the base; second cup-shaped; third and fourth very small; fifth and following joints to the tenth rather short, nearly equal in size; club spindle-shaped, more than twice the length of the tenth joint: chest elliptical, bluish green: fore-chest short; its length above little more than one-eighth of its breadth; hind-border slightly concave: sutures of the parapsides very indistinct; axillæ parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest of moderate size, obconical, declining: petiole very short: abdomen nearly linear, flat, smooth, purplish copper, bluish green at the base and at the tip, narrower but hardly shorter than the chest; metapodeon occupying about one-third of the back; octoon and following segments of moderate size: legs bright yellow; hips and thighs green; tips of feet piceous; fore-shanks and fore-feet tawny: wings colourless; veins dull tawny; ulna hardly half the length of the humerus; radius much longer than the ulna; cubitus shorter than the ulna; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Patro. Mas. *Cyaneo-viridis, abdomine æneo, macula viridi basali fulva, antennis fulvis, pedibus luteis, femoribus basi viridi vittatis, alis limpidis.* Page 125.

Bright green with a bluish tinge: head and chest convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers dark tawny, filiform, longer than the chest; first joint yellow, piceous at the tip; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club elliptical, more than twice the

length of the tenth joint ; each joint darker towards the base : chest oval, robust : fore-chest very short ; its length above about one-eighth of its breadth ; hind-border slightly concave : shield of the mid-chest broad ; sutures of the parapsides indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen bright coppery-green, smooth, flat, bronze on the disk, with a large tawny spot near the base, shorter and very much narrower than the chest ; metapodeon occupying about half of the back ; octoon short ; ennaton and following segments still shorter : legs luteous ; hips green ; each thigh with a green line near its base ; knees and feet of the four hinder-legs yellow ; tips of the feet brown : wings colourless ; veins dull tawny ; ulna about half the length of the humerus ; radius longer than the ulna ; cubitus much shorter than the ulna ; wing-brand piceous, small. Length of the body $1\frac{1}{4}$ line ; of the wings 2 lines.

Var. β . Hind-shanks yellow, with a tawny band across each.

PTEROMALUS LISSOS. MAS. *Viridis, abdominis disco aneo, antennis fulvis, pedibus flavis, alis limpidis.* Page 125.

Bright green : head and chest convex, finely shagreened : head a little broader than the chest : eyes and eyelets red : feelers tawny, subclavate, not longer than the chest ; first joint yellow, long, linear ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club brown, conical, more than twice the length of the tenth joint : chest oval : fore-chest very short ; its length less than one-tenth of its breadth ; hind-border slightly concave : shield of the middle-chest broad ; sutures of the parapsides very indistinct ; axillæ parted by one-third of the breadth of the chest ; scutcheon subrhomboidal : hind-chest short-obconical, slightly declining, with a slight ridge along the middle : petiole extremely short : abdomen nearly elliptical, smooth, flat, coppery on the disk, shorter and a little narrower than the chest ; metapodeon occupying nearly one-half of the back ; octoon and following segments of moderate size : legs bright yellow ; hips green ; tips of feet brown : wings colourless, rather large ; veins pale yellow ; ulna about half the length of the humerus ; radius longer than the ulna ; cubitus much shorter than the radius ; wing-brand very small. Length of the body $1\frac{1}{2}$ line ; of the wings $3\frac{1}{4}$ lines.

PTEROMALUS Morys. Mas. *Viridis, abdomine purpureo-cupreo, basi cyaneo-viridi, antennis fulvis, pedibus flavis, alis limpidis.* Page 125.

Bright green: head and chest convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers tawny, subclavate, a little longer than the chest; first joint bright yellow; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth short and successively decreasing in length; club elliptical, much more than twice the length of the tenth joint: chest nearly oval: fore-chest very short; its length above less than one-tenth of its breadth: shield of the mid-chest broad; sutures of the parapsides very indistinct; axillæ parted by nearly one-third of the breadth of the shield; scutcheon subrhomboidal: hind-chest of moderate size, short-obconical, declining, with a ridge along the middle: petiole very short: abdomen nearly linear, flat, smooth, purplish copper on the disk, bluish green at the base, a little narrower but hardly shorter than the chest; metapodeon occupying rather more than one-fourth of the back; octoon and ennaton rather large; the following segments very short: legs bright yellow; hips green; tips of feet brown: wings colourless; veins tawny; ulna brown, very thick, more than half the length of the humerus; radius longer than the ulna; cubitus more than half the length of the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Elyros. Mas. *Viridis, abdomine purpureo-cupreo, macula basali flava, antennis nigris, pedibus flavis, meta-femoribus fusco vittatis, alis limpidis.* Page 125.

Body long and narrow: head and chest convex, finely shagreened: head bluish green, much broader than the chest: eyes and eyelets red: feelers black, filiform, longer than the chest; first joint long, slender, piceous, yellow towards the base; second cup-shaped, piceous; third and fourth very small; the following joints from the fifth to the tenth long, linear, but successively decreasing in length; club linear, conical at the tip, more than twice the length of the tenth joint: chest green, spindle-shaped: fore-chest very short; its length above less than one-tenth of its breadth; hind-border slightly concave: shield of the mid-chest rather long; sutures of the parapsides very indistinct; axillæ parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a slight trace of a transverse ridge near its tip: hind-chest well developed, obconical, slightly declining, with a rim on each

side : petiole very short : abdomen smooth, flat, linear, conical at the base and pointed at the tip, purplish copper, with a pale yellow spot near the base which is green, shorter and narrower than the chest ; metapodeon occupying more than one-third of the back ; octoon and following segments of moderate size : legs yellow ; hips green ; a brown line along the whole length of each hind-thigh ; tips of feet brown : wings colourless ; veins tawny ; ulna nearly as long as the humerus ; radius shorter than the ulna ; cubitus very nearly half the length of the ulna ; wing-brand very small. Length of the body $1\frac{1}{4}$ line ; of the wings 2 lines.

PTEROMALUS Rhytium. Mas. *Viridis, abdominis disco purpureo-cupreo, antennis fuscis, pedibus flavis, alis limpidis.* Page 125.

Head and chest bright green, convex, finely shagreened : head much broader than the chest : eyes and eyelets red : feelers filiform, brown, rather hairy, a little longer than the chest ; first joint yellow, long, rather stout, piceous at the tip ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club spindle-shaped, more than twice the length of the tenth joint : chest elliptical : fore-chest very short ; its length less than one-tenth of its breadth : sutures of the parapsides very indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest rather large, obconical, declining, with a ridge along the middle : petiole very short : abdomen linear, flat, smooth, purplish copper, green at the base and at the tip, much narrower but hardly longer than the chest ; metapodeon occupying rather less than one-fourth of the back ; octoon and following segments of moderate size : legs bright yellow ; hips green ; middle-feet and hind-feet pale yellow with brown tips : wings colourless ; veins pale tawny ; ulna rather more than half the length of the humerus ; radius much shorter than the ulna ; cubitus a little shorter than the ulna ; wing-brand small. Length of the body 1 line ; of the wings $1\frac{1}{4}$ line.

Var. β . Chest coppery green : hind-chest bluish green.

PTEROMALUS Myle. Fem. *Æneus, abdomine viridi disco purpureo, antennis fulvis, pedibus rufis, alis fulvis.* Page 125.

Body short : head and chest convex, brassy, very finely shagreened : head large, much broader than the chest : eyes and eyelets dark red : feelers tawny, subclavate, shorter than the chest ; first joint pale red, long, slender ; second cup-shaped ; third and

fourth very small; the following joints from the fifth to the tenth short and successively decreasing in length; club elliptical, pointed at the tip, longer than the tenth joint and nearly thrice its length: chest elliptical: fore-chest extremely short; its length less than one-tenth of its breadth: shield of the middle-chest short; sutures of the parapsides very indistinct; axillæ parted by one-third of the breadth of the chest; scutcheon subrhomboidal: hind-chest short: petiole very short: abdomen short-elliptical, smooth, green, depressed above, very slightly keeled beneath, much shorter but hardly narrower than the chest; disk dark purple; metapodeon occupying full one-third of the back; the following segments successively decreasing in length: legs pale red; hips brassy; tips of feet piceous: wings narrow, tawny; veins dark tawny; ulna more than half the length of the humerus; radius a little shorter than the ulna; cubitus nearly as long as the radius; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{4}$ line.

PTEROMALUS Læta. Fem. *Viridis, abdomine purpureo basi cupreo, antennis fuscis, pedibus fulvis, femoribus aeneis, alis limpidis.* Page 125.

Body rather narrow: head and chest green, finely shagreened: head rather short, a little broader than the chest; crown and front broad: eyes and eyelets red, the former rather small: feelers brown, clavate, rather slender, much shorter than the chest; first joint tawny, slender, very long, hardly shorter than all the following joints together; second long cup-shaped; third and fourth very small; the following joints from the fifth to the tenth very short; club elliptical, pointed at the tip, and nearly thrice the length of the tenth joint: chest nearly linear, conical in front, flat above: fore-chest of moderate size, narrower in front; its breadth slightly exceeding its length: shield of the middle-chest broader than long; sutures of the parapsides very distinct on each side, but almost obsolete before they reach the hind-border; axillæ parted by one-third of the breadth of the chest; scutcheon purple, nearly truncate-conical, of moderate size, green along the hind-border: hind-chest rather large, short-obconical, nearly flat or very slightly declining, with an impression on each side of the fore-border: petiole very short: abdomen elliptical, smooth, purple, coppery at the base, depressed above, hardly keeled beneath, a little broader but not longer than the chest: legs tawny; hips and thighs brassy; mid-feet and hind-feet yellow with tawny tips: wings colourless, rather narrow and short; veins pale yellow; ulna more than half the length of

the humerus ; radius less than half the length of the ulna ; cubitus longer than the radius ; wing-brand very small. Length of the body 1 line ; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Androbius. Fem. *Viridis, abdomine viridi-purpureo, antennis nigris, pedibus fulvis femoribus piceis, alis limpidis.* Page 126.

Allied to *P. domesticus* but narrower : head and chest convex, dark green, finely shagreened : head broader than the chest : eyes and eyelets dark red : feelers black, clavate, not longer than the chest ; first joint tawny, long, slender ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth short and successively decreasing in length ; club conical, more than twice the length of the tenth joint : chest elliptical : fore-chest very short ; its length above less than one-tenth of its breadth : shield of the mid-chest short ; sutures of the parapsides very indistinct ; axillæ parted by almost one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen short-elliptical, smooth, purple, slightly tinged with green, bright green at the base, depressed above, hardly keeled beneath, a little broader but scarcely longer than the chest ; metapodeon occupying about one-sixth of the back ; octoon and following segments of moderate and nearly equal length : legs tawny ; hips brassy green ; thighs piceous ; middle-feet and hind-feet pale tawny with piceous tips : wings colourless ; veins tawny ; ulna hardly half the length of the humerus ; radius a little shorter than the ulna ; cubitus shorter than the radius ; wing-brand very small. Length of the body 1 line ; of the wings $1\frac{1}{4}$ line.

PTEROMALUS Inatos. Fem. *Cupreus, abdomine cupreo-purpureo, antennis nigris, pedibus fulvis, femoribus viridi-æneis, alis subfulvis.* Page 126.

Head and chest convex, coppery, finely shagreened : head a little broader than the chest : eyes piceous : feelers black, subclavate, shorter than the chest ; first joint long, tawny till near the tip ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club long-conical, more than twice the length of the tenth joint : chest short-oval : fore-chest very short ; its length above not more than one-tenth of its breadth ; hind-border slightly concave : su-

tures of the parapsides very indistinct; axillæ parted by rather less than one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest rather short, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen coppery-purple, smooth, nearly oval, rather slender towards the tip, depressed above, very much keeled beneath, rather broader and very much longer than the chest; a large triangular bright bluish green spot on each side of every segment; metapodeon golden green towards the base, occupying hardly more than one-eighth of the back: hind-border convex; octoon and ennaton of moderate size; decaton and the following segments rather longer: legs tawny; hips and thighs brassy green; mid-feet and hind-feet yellow with piceous tips: wings with a slight tawny tinge; veins dark tawny; ulna not more than half the length of the humerus; radius a little shorter than the ulna; cubitus much shorter than the radius; wing-brand piceous, very small. Length of the body 2 lines; of the wings 3 lines.

PTEROMALUS AXOS. Fem. Viridis, abdominis disco cupreo-purpureo, antennis piceis, pedibus flavis, alis limpidis. Page 126.

Bright green: head and chest convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers clavate, dark piceous, shorter than the chest; first joint tawny, long, slender; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth hardly decreasing in length; club nearly oval, pointed at the tip, a little broader than the tenth joint and more than twice its length: chest elliptical, with a bluish tinge: fore-chest very short; its length above not more than one-tenth of its breadth; hind-border slightly concave: sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side, and a very indistinct transverse suture near the tip: hind-chest of moderate size, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen long-oval, smooth, depressed above, very slightly keeled beneath, a little broader but hardly longer than the chest; disk coppery purple, which colour spreads more along the hind-border than along the fore-border of each segment; metapodeon not occupying more than one-sixth of the back; hind-border slightly concave; all the following segments of moderate and nearly equal length: legs yellow; hips green; tips of feet brown; middle-feet and hind-feet pale yellow: wings colourless; veins tawny; ulna full half the

length of the humerus ; radius a little shorter than the ulna ; cubitus much shorter than the radius ; wing-brand small, brown. Length of the body $1\frac{1}{4}$ line ; of the wings 2 lines.

PTEROMALUS Otos. Fem. *Viridis, abdomine cupreo-purpureo, antennis nigris, pedibus fulvis, femoribus viridibus, alis subfulvis.* Page 126.

Head and chest dark green, convex, finely shagreened : head large, broader than the chest : eyes and eyelets dark red : feelers black, clavate, nearly as long as the chest ; first joint long, slender, tawny, black towards the tip ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth short and successively decreasing in length ; club short, nearly conical, broader than the tenth joint and about twice its length : chest oval : fore-chest very short ; its length not more than one-tenth of its breadth ; sutures of the parapsides very indistinct ; axillæ parted by nearly one-fourth of the breadth of the chest ; scutcheon subrhomboidal, having a suture along each side, and a slight trace of a cross furrow near its tip : hind-chest rather large, obconical, slightly declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen oval, smooth, coppery-purple, bright green at the base, flat above, keeled beneath, shorter and a little broader than the chest ; metapodeon occupying full one-fourth of the back ; octoon and all the following segments of moderate and nearly equal size : legs tawny ; hips and thighs green ; tips of the latter tawny ; middle-feet and hind-feet yellow with brown tips : wings with a slight tawny tinge ; veins tawny ; ulna about half the length of the humerus ; radius a little shorter than the ulna ; cubitus shorter than the radius ; wing-brand very small. Length of the body $1\frac{1}{4}$ line ; of the wings 2 lines.

PTEROMALUS Carina. Fem. *Viridis, abdominis disco purpureo, antennis fulvis, pedibus flavo-fulvis, femoribus basi piceis, alis limpidis.* Page 126.

Bright green : head and chest convex, finely shagreened : head a little broader than the chest : eyes and eyelets red : feelers dark tawny, filiform, hardly longer than the chest ; first joint long, slender, yellow, piceous towards the tip ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth equal in size and rather broader than long ; club spindle-shaped, thrice the length of the tenth joint : chest nearly elliptical : fore-chest very short ; its length above less than one-tenth of its

breadth ; hind-border very slightly concave : shield of the mid-chest rather short ; sutures of the parapsides very indistinct ; axillæ parted by about one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest short, slightly obconical, declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen oval, smooth, depressed above, hardly keeled beneath, scarcely longer or broader than the chest ; disk purple, which colour mingled with a brassy tinge extends along the hind-border of each segment ; metapodeon occupying full one-sixth of the back ; hind-border slightly convex ; octoon and following segments of moderate and equal size : legs bright pale tawny ; hips green ; thighs piceous towards the base ; four hinder-knees and feet pale yellow, tips of the latter piceous : wings colourless ; veins pale tawny ; ulna hardly half the length of the humerus ; radius a little longer than the ulna ; cubitus much shorter than the ulna ; wing-brand pale brown, very small. Length of the body 1 line ; of the wings 2 lines.

PTEROMALUS Sybritia. Fem. Viridis, cyaneo cupreo purpureoque varius, antennis nigris, pedibus flavo-fulvis, femoribus piceo-vittatis, alis limpidis. Page 126.

Head and chest convex, finely shagreened : green slightly tinged with blue and copper colour : head large, broader than the chest : eyes and eyelets dark red : feelers black, subclavate, rather slender, piceous towards the tip ; first joint long, slender, tawny towards the base ; second long-cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club long-conical, more than twice the length of the tenth joint : chest elliptical : fore-chest very short ; its length above less than one-tenth of its breadth ; hind-border slightly concave : sutures of the parapsides indistinct ; axillæ parted by rather less than one-fourth of the breadth of the chest ; scutcheon subrhomboidal, with a suture along each side : hind-chest rather short, declining, golden green, almost smooth, with an impression on each side of the fore-border : petiole very short : abdomen long-oval, smooth, bright bluish green, depressed above, keeled beneath, very little narrower and shorter than the chest ; disk purple ; angle of the keel very near the base ; metapodeon occupying about one-fourth of the back ; octoon and three following segments of moderate size ; paratelum and telum rather short : legs bright tawny ; hips green ; thighs with piceous stripes ; four hinder-shanks and feet bright pale yellow ; tips of the latter piceous : wings colourless ; veins pale tawny ; ulna about half the length of the humerus ; ra-

dus nearly as long as the ulna ; cubitus much shorter than the radius ; wing-brand very small. Length of the body $1\frac{1}{4}$ line ; of the wings $2\frac{1}{2}$ lines.

PTEROMALUS Mesapos. Fem. *Cupreus, abdomine cupreo-purpureo basi viridi, antennis nigris, pedibus flavis, femoribus fulvis piceo vittatis, alis limpidis.* Page 126.

Head and chest coppery, convex, finely shagreened : head broader than the chest : eyes and eyelets dark red : feelers black, clavate, thick, a little longer than the chest ; first joint bright tawny, long, linear, black at the tip ; second cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club conical, much broader than the tenth joint and more than twice its length : chest nearly oval : fore-chest very short ; its length above less than one-tenth of its breadth ; hind-border very concave : sutures of the parapsides very indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest rather large, obconical, declining, with some excavations along the fore-border and a slight ridge along the middle : petiole very short : abdomen oval, smooth, coppery purple, bright green at the base, depressed above, very deeply keeled beneath, so as almost to form a right angle, shorter and a little narrower than the chest ; metapodeon occupying about one-fourth of the back, slightly excavated at the base ; octoon and the following segments of moderate and nearly equal size ; parate-lum and telum rather long ; legs yellow, rather stout, hips brassy ; thighs tawny, with piceous stripes ; each middle-thigh with a spine beneath near the tip ; four hinder-feet with piceous tips : wings colourless, rather short ; veins pale tawny ; ulna more than half the length of the humerus ; radius a little shorter than the ulna ; cubitus much shorter than the radius ; wing-brand very small, pale brown. Length of the body $1\frac{1}{4}$ line ; of the wings 2 lines.

PTEROMALUS Larymna. Fem. *Viridis, abdomine viridi-cyaneo, antennis nigris, pedibus fulvo-piceis, femoribus viridibus, alis limpidis.* Page 126.

Bright green : head and chest convex, shagreened : head hardly as broad as the chest : eyes and eyelets bright red : feelers black, clavate, slender, much shorter than the chest ; first joint long, green, slightly increasing in breadth from the base to the tip ; second long cup-shaped ; third and fourth very small ; the following joints from the fifth to the tenth short and successively decreasing

in length; club oval, much broader than the tenth joint and more than twice its length; chest elliptical: fore-chest very short; its length above about one-eighth of its breadth; hind-border very slightly concave: shield of the middle-chest rather long; sutures of the parapsides rather indistinct on each side and almost obsolete before they reach the hind-border; axillæ parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest short, transverse: petiole very short: abdomen oval, smooth, greenish blue, depressed above, hardly keeled beneath, a little longer and broader than the chest; metapodeon occupying about one-sixth of the back; octoon and following segments of moderate and nearly equal length: hips and thighs green; shanks and tips of four hinder-feet piceous; knees, feet, and tips of shanks tawny: wings colourless; veins brown; ulna less than half the length of the humerus; radius much longer than the ulna; cubitus shorter than the ulna; wing-brand small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{2}$ lines.

PTEROMALUS Etearchus. Fem. *Viridis, abdomine purpureo-viridi, antennis nigris, pedibus flavo-luteis, alis subfulvis.*
Page 126.

Body green: head and chest convex, finely shagreened: head broader than the chest: eyes and eyelets red: feelers black, subclavate, rather slender, not longer than the chest; first joint long, slender, tawny, black at the tip; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club oval, much broader than the tenth joint and more than twice its length: chest elliptical: fore-chest very short; its length less than one-tenth of its breadth; hind-border slightly concave: shield of the middle-chest rather long; sutures of the parapsides very indistinct; axillæ parted by rather less than one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest rather short, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, smooth, depressed above, keeled and tawny beneath, narrower and very much longer than the chest; disk tinged with purple; metapodeon occupying about one-sixth of the back; octoon and three following segments of moderate and nearly equal size; paratelum and telum shorter: legs luteous; four hinder-shanks and feet bright yellow; tips of the latter brown: wings with a tawny tinge; veins tawny; ulna rather broad, about half the length of the humerus; radius a little longer than the ulna; cubitus much shorter than the ulna; wing-brand small. Length of the body $1\frac{1}{4}$ line; of the wings $1\frac{3}{4}$ line.

PTEROMALUS Phalarsaina. Fem. *Viridis, abdomine purpureo-cupreo, basi viridi, antennis nigris, pedibus fulvis, femoribus anticis piceo-vittatis, alis subfuscis.* Page 126.

Allied to *Trigonoderus*. Body rather narrow : head and chest green, convex, finely shagreened : head rather large, broader than the chest : eyes and eyelets dark red : feelers black, subclavate, nearly as long as the chest ; first joint green, long, slender ; second cup-shaped ; third and fourth very short ; the following joints, from the fifth to the tenth successively, decreasing in length ; club long-conical, more than twice the length of the tenth joint : chest spindle-shaped, well developed : fore-chest rather large ; its length about one-fourth of its breadth ; hind border concave : shield of the mid-chest narrower in front : sutures of the parapsides distinct ; axillæ parted by about one-sixth of the breadth of the chest ; scutcheon subrhomboidal, nearly conical in front, with a suture along each side and a transverse suture near the tip : hind-chest well developed, obconical, declining, roughly sculptured : fore-shield distinct : petiole very short : abdomen spindle-shaped, smooth, purplish copper, green at the base, depressed above, slightly keeled beneath, very little shorter and narrower than the chest ; metapodeon occupying about one-fourth of the back ; octoon and all the following segments of moderate and nearly equal size : legs tawny ; hips green ; fore-thighs striped with green ; four hinder feet with piceous tips : wings slightly tinged with brown ; veins dull tawny ; ulna a little more than half the length of the humerus ; radius a little longer than the ulna ; cubitus much shorter than the ulna ; wing-brand very small. Length of the body $1-1\frac{1}{2}$ line ; of the wings $2-2\frac{1}{2}$ lines.

Var. β . Thighs green ; four hinder feet much paler than the shanks.

PTEROMALUS Themiso. Fem. *Viridis, abdomine cupreo-purpureo, basi viridi micante, antennis nigris, pedibus fulvis, femoribus viridi vittatis, alis fulvis.* Page 126.

Body rather narrow : head and chest dark green, convex, finely shagreened : head rather large, broader than the chest : eyes and eyelets dark red : feelers black, clavate, rather thick, nearly as long as the chest ; first joint, long, linear, tawny, piceous at the tip ; second cup-shaped ; third and fourth very short ; the following joints to the tenth successively decreasing in length ; club conical, hardly broader than the tenth joint, but more than twice its length : chest spindle-shaped : fore-chest of moderate size ; its length about one-fourth of its breadth : shield of the mid-chest rather long : sutures of the parapsides very indistinct ; axillæ parted by rather less

than one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest large, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, smooth, coppery purple, bright green at the base, depressed above, keeled beneath, as long and as broad as the chest; metapodeon occupying nearly one-fourth of the back; octoon and three following segments tolerably long, nearly equal in size; paratelum and telum short: legs tawny; hips green; thighs striped with green; four hinder-knees and feet yellow, tips of the latter tawny: wings tawny, rather narrow; veins of the same colour; ulna more than half the length of the humerus; radius a little shorter than the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body 1 line; of the wings $1\frac{3}{4}$ line.

PTEROMALUS Allutius. Fem. Viridis, cyaneo cupreo purpureoque varius, antennis nigris, pedibus fulvis, alis subfulvis.
Page 126.

Body rather short: head and chest convex, finely shagreened: head green, a little broader than the chest: eyes and eyelets red: feelers black, clavate, a little shorter than the chest; first joint tawny, long, and very slender; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club oval, much broader than the tenth joint and more than twice its length: chest elliptical, purplish copper, not much developed: fore-chest very short; its length less than one-tenth of its breadth; hind-border slightly concave: shield of the mid-chest short; sutures of the parapsides very indistinct, almost obsolete; axillæ parted by one-third of the breadth of the chest; scutcheon subrhomboidal, with a short suture along each side: petiole very short: abdomen oval, smooth, varied with blue, purple, green, and copper-colour, flat above, slightly keeled beneath, broader but scarcely longer than the chest; metapodeon occupying about one-fifth of the back; octoon and all the following segments of moderate and nearly equal length: legs tawny; hips coppery; four hinder-knees and feet yellow; tips of the feet brown: wings rather small, with a very slight tawny tinge; veins pale yellow; ulna less than half the length of the humerus; radius a little longer than the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{4}$ line.

PTEROMALUS Priansos. Fem. *Viridi-cyaneus, abdominis disco nigro purpureo, antennis pedibusque fulvis, femoribus cyaneo-viridibus, tibiis piceo fasciatis, alis limpidis.* Page 126.

Greenish blue: head and chest convex, finely shagreened: head as broad as the chest: eyes and eyelets red: feelers piceous, clavate, slender, rather shorter than the chest; first joint long, slender, tawny at the base; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club oval, much broader than the tenth joint and more than twice its length: chest short-elliptical: fore-chest short; its length about one-eighth of its breadth: shield of the mid-chest rather long; sutures of the parapsides indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon sub-rhomboidal, with a suture along each side: hind-chest transverse, short, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, smooth, depressed above, deeply keeled beneath, narrower and much longer than the chest; disk blackish purple; metapodeon occupying rather less than one-fourth of the back; octoon and ennaton of equal size; deca-ton and protelum longer; paratelum and telum shorter: legs tawny; hips and thighs bluish green; a broad piceous band across each of the four hinder-shanks; four hinder-feet pale tawny with piceous tips: wings colourless; veins pale tawny; ulna hardly half the length of the humerus; radius about as long as the ulna; cubitus much shorter than the radius: wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

Var. β . Head and chest green.

PTEROMALUS Gigon. Fem. *Viridis, abdomine purpureo-æneo, antennis nigris, pedibus fulvis, alis limpidis.* Page 126.

Allied to *P. domesticus*. Body rather short: head and chest green, convex, finely shagreened: head very little broader than the chest: eyes and eyelets red: feelers black, clavate, shorter than the chest; first joint green, long, slender; second cup-shaped; third and fourth very small; the following joints to the tenth short, transverse; club conical, hardly broader than the tenth joint but more than twice its length: chest elliptical, not much developed: fore-chest extremely short; its length above much less than one-tenth of its breadth: shield of the mid-chest short; sutures of the parapsides very indistinct, almost obsolete; axillæ parted by full one-fourth of the chest; scutcheon subrhomboidal: hind-chest transverse, rather short, slightly declining, with a slight ridge along the middle and a

rim on each side: petiole very short: abdomen elliptical, smooth (that is, comparatively, for in this and several other species of *Pteromalus*, it appears slightly shagreened when highly magnified), dark purplish bronze, brassy at the base and at the tip, depressed above, hardly keeled beneath, very little longer and broader than the chest; metapodeon as long as the octoon, hind-border slightly convex with anotch in the middle; octoon and four following segments of moderate and nearly equal size; telum very small: legs tawny; hips green; tips of the feet brown: wings colourless, rather short; veins pale tawny; ulna rather less than half the length of the humerus; radius a little shorter than the ulna; cubitus shorter than the radius; wing-brand extremely small. Length of the body 1 line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Orthagus. Fem. Cyaneo-viridis, abdomine cupreo-purpureo, antennis nigris, tibiis piceis, tarsis fulvis, alis limpidis. Page 126.

Head and chest bluish green, convex, finely shagreened: head as broad as the chest: eyes and eyelets red: feelers black, shorter and more clavate than those of *P. Priansos*; first joint green, long, slender; second cup-shaped; third and fourth very small; the following joints to the tenth short, transverse; club nearly oval, broader than the tenth joint and more than twice its length: chest oval: fore-chest very short; its length not more than one-tenth of its breadth: shield of the middle-chest rather large; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest transverse, rather short, with a slight ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, smooth, blue, coppery purple on the disk, bright green at the base, depressed above, slightly keeled beneath, longer and much narrower than the chest; most of the segments of moderate and nearly equal length: hips and thighs green; shanks piceous; knees and feet tawny; tips of four hinder-feet piceous: wings colourless; veins pale tawny; ulna hardly half the length of the humerus; radius as long as the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings 2 lines.

PTEROMALUS Diomedon. Fem. Purpureo-viridis, abdomine viridi-cyaneo, antennis nigris, pedibus flavo-piceis, femoribus viridibus, alis limpidis. Page 126.

Head and chest dark green tinged with purple: head very little

broadier than the chest: eyes and eyelets dark red: feelers black, subclavate, rather slender, a little longer than the chest; first joint green, long, linear; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club long-oval, a little broader than the tenth joint and more than twice its length: chest short-elliptical: fore-chest very short; its length above less than one-tenth of its breadth: shield of the mid-chest rather short; sutures of the parapsides indistinct; axillæ parted by little more than one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest golden green, short, transverse, with a slight ridge along the middle and a rim on each side: petiole very short: abdomen spindle-shaped, smooth, greenish blue, depressed above, keeled beneath, a little narrower and very much longer than the chest; metapodeon occupying nearly one-fifth of the back; octoon short; ennaton and following segments longer: hips and thighs green; shanks piceous; fore-knees and fore-feet tawny; knees, feet, and tips of shanks of four hinder-legs yellow: wings colourless; veins pale yellow; ulna about half the length of the humerus; radius as long as the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body 1 line; of the wings 2 lines.

PTEROMALUS Alimentus. Fem. Nigro-æneus, abdomine purpureo basi viridi, antennis femoribusque, piceis pedibus flavis, alis subfulvis. Page 126.

Head and chest convex, dull black, with a very slight brassy tinge, finely shagreened: head hardly broader than the chest: eyes and eyelets dark red: feelers dark piceous, subclavate, not longer than the chest; first joint long, slender, yellow, piceous at the tip; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest nearly oval: fore-chest very short: shield of the middle-chest nearly flat; sutures of the parapsides very indistinct; axillæ parted by nearly one-third of the breadth of the chest; scutcheon subrhomboidal: hind-chest short, hardly obconical, declining: petiole very short: abdomen spindle-shaped, smooth, dark purple, green at the base, depressed above, keeled beneath, narrower and very much longer than the chest; metapodeon occupying about one-fifth of the back; octoon and following segments of moderate size and of nearly equal length: legs bright yellow; hips black; thighs except the tips, piceous; middle-feet and hind-feet pale yellow with

black tips: wings pale tawny; a large brown spot on the disk of each fore-wing; veins pale brown; ulna hardly half the length of the humerus; radius a little longer than the ulna; cubitus much shorter than the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ —1 line; of the wings $1\frac{1}{4}$ — $1\frac{1}{2}$ line.

Var. β . Feelers black.

PTEROMALUS Cælius. Fem. *Æneus, abdomine purpureo, antennis nigris, pedibus fulvis, femoribus piceo-vittatis, proalis fusco submaculatis*. Page 126.

Brassy: head and chest convex, shagreened: head a little broader than the chest: eyes and eyelets red: feelers black, almost filiform or very slightly clavate, not longer than the chest; first joint tawny, long, slender; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club spindle-shaped, more than twice the length of the tenth joint: chest oval: fore-chest extremely short: sutures of the parapsides tolerably distinct; axillæ parted by about one-fifth of the breadth of the chest; scutcheon subrhomboidal with a transverse suture near its tip: hind-chest rather large, short-obconical, declining, with a ridge along the middle and a rim on each side; its fore-shield distinct: petiole extremely short: abdomen spindle-shaped, smooth, dark purple, depressed on the disk, hardly keeled beneath, a little narrower and much longer than the chest; metapodeon occupying about one-sixth of the back, notched in the middle of the hind-border; octoon like the following segments of moderate size, and nearly equal to the metapodeon in length: sheaths of the oviduct black, hairy: legs pale tawny; hips brassy; a piceous line along the upper border of each of the fore-thighs and of the hind-thighs; tips of the feet brown: wings colourless; each fore-wing with a brown spot beneath the brand; veins tawny; ulna about half the length of the humerus; radius very much longer than the ulna; cubitus more than half the length of the ulna; wing-brand small. Length of the body 2 lines; of the wings 3 lines.

PTEROMALUS Æacus. Fem. *Cyaneo-viridis, abdomine purpureo nigro, antennis piceis, pedibus piceo-fulvis, alis limpidis*. Page 126.

Head and chest convex, finely shagreened: head dark blue with a purple tinge, not broader than the chest: eyes and eyelets dark red: feelers piceous, clavate, not longer than the chest; first

joint long, slender, green; second cup-shaped; third and fourth very small; fifth and following joints very short; club oval, much broader than the tenth joint and more than twice its length: chest oval, dark green with a blue tinge: fore-chest very short; its length above not more than one-tenth of its breadth; shield of the mid-chest flat; sutures of the parapsides very indistinct; axillæ parted by about one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest of moderate size, obconical, declining, with a suture on each side: petiole piceous, very short: abdomen nearly oval, smooth, purplish black, dark bluish green at the base, depressed above, slightly keeled beneath, a little shorter and narrower than the chest; metapodeon occupying less than half of the back; octoon and two following segments of moderate size; the other segments very short: legs pale tawny; hips dark green; thighs and shanks of four hinder-feet slightly piceous; middle-feet and hind-feet yellow with brown tips: wings colourless; veins pale tawny; ulna full half the length of the humerus; radius longer than the ulna; cubitus much shorter than the ulna; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Alope. Fem. Purpureo-niger, abdomine viridivario, antennis nigris, pedibus piceis, alis subfuscis. Page 127.

Head and chest convex, finely shagreened, black with a slight purple tinge: head not broader than the chest: eyes and eyelets piceous: feelers black: chest short-elliptical: fore-chest very short; its length above not more than one-tenth of its breadth; hind-border hardly concave: sutures of the parapsides indistinct; axillæ parted by nearly one-fifth of the breadth of the chest; scutcheon subrhomboidal: hind-chest of moderate size, nearly obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen nearly elliptical, smooth, dark purple slightly tinged with green, depressed above, keeled beneath, a little broader but not longer than the chest; metapodeon nearly half the length of the back; octoon and following segments short: legs piceous; hips black: wings pale brown; veins pale brown; ulna not more than half the length of the humerus; radius longer than the ulna; cubitus more than half the length of the ulna; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Cepio. Fem. *Cyaneo-viridis, abdominis disco purpureo, antennis piceis, pedibus fulvis, alis limpidis.* Page 127.

Dark green : head and chest convex, finely shagreened : head dark bluish green, broader than the chest ; feelers piceous ; first joint tawny, slender, very long ; second darker, cup-shaped ; third and fourth very small ; fifth and the following joints to the tenth transverse, very short ; club long-conical, more than twice the length of the tenth joint : chest nearly elliptical : fore-chest very short ; its length above about one-eighth of its breadth ; hind-border concave : shield of the mid-chest rather flat ; sutures of the parapsides very indistinct ; axillæ parted by about one-fourth of the breadth of the chest ; scutcheon subrhomboidal : hind-chest rather short, slightly obconical, declining, with a ridge on each side : petiole very short : abdomen smooth, dark green, dark purple on the disk, depressed above, slightly keeled beneath, a little narrower and shorter than the chest ; metapodeon occupying about one-third of the back ; octoon and following segments of moderate size : legs pale tawny ; hips dark green ; thighs darker towards the base ; middle-feet and hind-feet yellow with tawny tips : wings colourless ; veins pale tawny ; ulna about half the length of the humerus ; radius not longer than the ulna ; cubitus much shorter than the radius ; wing-brand very small. Length of the body $\frac{2}{3}$ line ; of the wings $1\frac{1}{4}$ line.

PTEROMALUS Hyrtacina. Fem. *Viridis, abdominis disco cupreo-purpureo, antennis pedibusque fulvis, metafemoribus fusco vittatis, alis limpidis.* Page 127.

Green : head and chest convex, finely shagreened : head broader than the chest : eyes and eyelets red : feelers tawny, subclavate, rather longer than the chest ; first joint long, slender, tawny, piceous at the tip ; second cup-shaped, piceous ; third and fourth very small ; the following joints from the fifth to the tenth successively decreasing in length ; club spindle-shaped, rather broader than the tenth joint and about thrice its length : chest nearly oval, rather short : fore-chest very short ; its length less than one-tenth of its breadth : shield of the mid-chest broad ; sutures of the parapsides very indistinct ; axillæ parted by one-fourth of the breadth of the chest ; scutcheon subrhomboidal, with a suture along each side : hind-chest transverse, rather short, with a slight ridge along the middle and a rim on each side : petiole very short : abdomen spindle-shaped, smooth, depressed above, slightly keeled

beneath, longer and much narrower than the chest; disk coppery purple; metapodeon occupying about one-fourth of the back; ootoon and following segments of moderate length; legs pale tawny; hips green; hind-thighs striped with brown; four hinder-knees and feet yellow, tips of the latter brown; wings colourless, ample; veins pale tawny; ulna hardly half the length of the humerus; radius a little longer than the ulna; cubitus much shorter than the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

PTEROMALUS Leuce. Fem. *Æneo-viridis, abdominis disco purpureo, antennis fulvis, pedibus fulvo-flavis, alis limpidis.* Page 127.

Body nearly linear, brassy green: head and chest convex, finely shagreened: head as broad as the chest: eyes and eyelets red: feelers tawny, clavate, a little longer than the chest; first joint long, linear, yellow at the base; second cup-shaped; third and fourth very small; the following joints from the fifth to the tenth successively decreasing in length; club nearly oval, much broader than the tenth joint and more than twice its length: chest short-elliptical: fore-chest very short; its length above less than one-tenth of its breadth: sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest transverse, short: petiole very short: abdomen long-oval, smooth, depressed above, keeled beneath, much longer but not narrower than the chest; disk dark purple; metapodeon and all the following segments of moderate and nearly equal size: legs very pale tawny; hips green; four hinder-shanks and feet yellow, tips of the latter tawny: wings colourless, rather narrow; veins very pale yellow; ulna about half the length of the humerus; radius longer than the ulna; cubitus much shorter than the ulna; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{4}$ line.

PTEROMALUS Hyrtacus. Fem. *Viridi-aeneus, abdomine purpureo, antennis nigris, pedibus fulvis, femoribus piceo-vittatis, alis subfulvis.* Page 127.

Head and chest convex, finely shagreened: head very dark green, a little broader than the chest: eyes and eyelets red: feelers black, slightly subclavate, a little longer than the chest; first joint tawny, long, slender, linear; second cup-shaped, piceous; third and fourth very small; the following joints from the fifth to

the tenth successively decreasing in length; club long-conical, a little more than twice the length of the tenth joint: chest brassy, nearly elliptical: fore-chest very short; its length above not more than one-tenth of its breadth; hind-border concave: shield of the mid-chest narrower in front; sutures of the parapsides indistinct; axillæ parted by about one-fifth of the breadth of the chest; scutcheon subrhomboidal: hind-chest rather long, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen almost spindle-shaped, depressed above, very slightly keeled beneath, smooth, coppery purple, slender towards the tip, tawny beneath and at the base above, a little longer but hardly narrower than the chest; metapodeon tinged with green, occupying about one-fourth of the back; octoon and all the following segments of moderate and nearly equal length; telum slender and pointed: legs tawny; hips brassy; a brown streak along each of the four hinder-thighs; middle-feet and hind-feet yellow with brown tips: wings slightly tawny, rather narrow; veins brown; ulna a little more than half the length of the humerus; radius longer than the ulna; cubitus rather long, about half the length of the radius; wing-brand very small. Allied to *Hetroxys*. Length of the body $1\frac{1}{2}$ line; of the wings 2 lines.

Var. β . Chest green.

HETROXYS Aponius. Mas. *Æneo-viridis, abdomine purpureo-cupreo fulvo maculato, antennis pedibusque fulvis, femoribus piceis, alis limpidis*. Fem. *Abdomine immaculato, basi cyaneo-viridi, antennis nigris*. Page 127.

Male. Body slender: head and chest convex, shagreened: head dark green, a little broader than the chest: eyes and eyelets red: feelers dark tawny, slender, subclavate, much longer than the chest; first joint long, slender, tawny; second cup-shaped; third and fourth very small; the following joints to the tenth of equal breadth, but successively decreasing in length; club long-conical, a little broader than the tenth joint and nearly twice its length: chest spindle-shaped, dark coppery green: fore-chest very short; its length above not one-tenth of its breadth: shield of the mid-chest long, rather flat above; sutures of the parapsides very indistinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side, and a very indistinct transverse channel near the tip: hind-chest well developed, obconical, declining, with a ridge along the middle and a rim on each side: petiole very short: abdomen depressed, smooth, dark purplish copper, with a tawny spot near the base, slightly increasing in

breadth from the base till near the tip, shorter and much narrower than the chest; metapodeon not occupying more than one-sixth of the back; octoon and following segments of nearly equal size: legs tawny; hips green; thighs piceous towards the base; four hinder feet yellow with brown tips: wings colourless; veins tawny; ulna more than half the length of the humerus; radius much longer than the ulna; cubitus much shorter than the ulna; wing-brand large.

Female. Feelers black, a little longer than the chest; first joint tawny; abdomen long spindle-shaped, smooth, bluish green, purplish bronze on the disk, concave above, slightly keeled beneath, much longer and narrower than the chest, hairy towards the tip; metapodeon not occupying more than one-eighth of the back: octoon and ennaton short; decaton and three following segments longer. Length of the body $1-1\frac{1}{2}$ line; of the wings $2-2\frac{1}{2}$ lines.

TRIGONODERUS Lappa. Mas. *Cyaneo-viridis*, abdomine *purpureo-cupreo*, *antennis nigris*, *pedibus piceis*, *tarsis flavis*, *alis limpidis*. Page 128.

Body narrow: head and chest convex, finely shagreened, dark bluish green: head much broader than the chest: eyes and eyelets dark red: feelers black, filiform, rather broad, shorter than the chest; first joint long, spindle-shaped; second cup-shaped; third and fourth very short; the following joints to the tenth successively decreasing in length; club spindle-shaped, rather more than twice the length of the tenth joint: chest long-elliptical: fore-chest long; its length more than one-half of its breadth: shield of the mid-chest short and rather flat; sutures of the parapsides strongly marked; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, rather flat, with a suture along each side, and a slight transverse channel near the tip: hind-chest large, obconical, slightly declining, with no ridge nor rim: petiole very short: abdomen depressed, smooth, dark purplish bronze-colour, slightly increasing in breadth from the base till near the tip, much shorter and narrower than the chest; metapodeon long: legs piceous; hips green; fore-feet pale tawny; four hinder-feet yellow with brown tips: wings colourless, rather short; veins pale tawny; ulna full half the length of the humerus; radius much longer than the ulna; cubitus a little more than half the length of the ulna; wing-brand very small. Length of the body $\frac{3}{4}-1$ line; of the wings $1\frac{1}{4}-1\frac{1}{2}$ line.

Var. *β. Aut species distincta.* Thighs and shanks dull tawny; veins of the wings dark tawny.

TRIGONODERUS Tarrha. Mas. *Cyaneo-viridis, abdomine purpureo-cupreo basi viridi, antennis nigris, pedibus fulvis, femoribus piceo vittatis, alis limpidis.* Page 128.

Body narrow : head and chest convex, finely shagreened : head bright blue, much broader than the chest : eyes and eyelets red : feelers black, slender, hairy, very slightly subclavate, a little longer than the chest ; first joint long, slender, dark tawny ; second cup-shaped ; third and fourth very small ; the following segments to the tenth successively decreasing in length ; club spindle-shaped, a little broader than the tenth joint, and rather more than twice its length : chest spindle-shaped, bluish green : fore-chest rather long, narrower in front ; its length above more than half its breadth : shield of the mid-chest long ; sutures of the parapsides strongly marked ; axillæ parted by one-sixth of the breadth of the chest ; scutcheon subrhomboidal, with a suture along each side, and a slight transverse channel near the tip : hind-chest large, obconical, declining, with a ridge along the middle and a rim on each side : petiole very short : abdomen depressed, smooth, purplish copper, green at the base, very slightly increasing in breadth from the base till near the tip, shorter and a little narrower than the chest ; metapodeon occupying rather more than one-fourth of the back ; octoon and following segments of nearly equal length : legs pale tawny ; hips green ; a piceous line along each thigh ; four hinder-feet yellow with brown tips : wings colourless ; veins piceous ; ulna much more than half the length of the humerus ; radius a little longer than the ulna, reaching almost to the tip of the wing ; cubitus not half the length of the ulna ; wing-brand very small. Length of the body $1\frac{1}{4}$ line ; of the wings 2 lines.

TRIGONODERUS Alebion. Mas. *Viridis, capite cyaneo, abdominis, disco purpureo-cupreo, antennis nigris, pedibus piceo-fulvis femoribus, viridibus, alis limpidis.* Page 128.

Head and chest convex, shagreened : head blue, broader than the chest : eyes and eyelets red : feelers black, subclavate, a little longer than the chest ; first joint long, slender, green ; second cup-shaped ; third and fourth very small ; the following joints to the tenth of equal thickness but successively decreasing in length ; club spindle-shaped, a little broader than the tenth joint and nearly twice its length : chest green, spindle-shaped : fore-chest rather long ; its length more than half its breadth : shield of the mid-chest long ; sutures of the parapsides rather distinct ; axillæ parted by hardly one-sixth of the breadth of the chest ; scutcheon subrhomboidal, with a

suture along each side, and a slight transverse channel near the tip : hind-chest large, obconical, declining, with a slight ridge along the middle and a rim on each side, and having also a transverse ridge near the tip : petiole very short : abdomen depressed, smooth, nearly linear or rather narrower at the base than at the tip, green, dark purplish copper-colour on the disk, much narrower than the chest, and hardly more than half its length ; metapodeon occupying about half the back ; octoon and following segments successively decreasing in length : legs tawny ; hips and thighs green ; four hinder legs with yellow knees and feet, and with piceous shanks and tips of feet : humerus colourless ; veins piceous ; ulna hardly half the length of the humerus ; radius much longer than the ulna ; cubitus more than half the length of the ulna ; wing-brand very small. Length of the body $1\frac{1}{2}$ line ; of the wings $\frac{1}{2}$ line.

TRIGONODERUS Polichna. Mas. *Viridi-cyaneus, antennis nigris pedibus flavis, mesotibiis fulvis, alis limpidis.* Page 128.

Head and chest convex, finely shagreened, very dark bluish green : head much broader than the chest : eyes and eyelets dark red : feelers black, slender, subelavate, hairy, a little shorter than the body ; first joint long, slender ; second cup-shaped ; third and fourth very small ; the following joints to the tenth long, but successively decreasing in length ; club spindle-shaped, a little broader than the tenth joint, and nearly twice its length : chest long, spindle-shaped : fore-chest narrow and declining in front ; its length above about one-eighth of its breadth : shield of the mid-chest long ; sutures of the parapsides indistinct ; axillæ parted by rather less than one-fourth of the breadth of the chest ; scutcheon sub-rhomboidal, with a suture along each side, and a transverse indistinct channel near the tip : hind-chest large, obconical, declining, with no ridge nor rim : petiole very short : abdomen depressed, finely shagreened, bluish green, very slightly increasing in breadth from the base till near the tip, narrower but not shorter than the chest ; metapodeon occupying about one-fifth of the back ; octoon and following segments rather shorter and of equal length : legs yellow ; hips green ; fore-shanks and fore-feet dull yellow ; middle shanks and tips of four hinder feet dark tawny : wings colourless ; veins tawny ; ulna rather stout, a little more than half the length of the humerus ; radius a little longer than the ulna ; cubitus not half the length of the ulna ; wing-brand very small. Length of the body $1\frac{1}{2}$ line ; of the wings 2 lines.

EUPELMUS ACRON. Fem. *Æneo-viridis, capite purpureo, antennis nigris, pedibus fulvis, femoribus tibiisque anticis piceis, alis fulvis.* Page 129.

Connects Eupelmus with Trigonoderus. Head finely shagreened, nearly smooth, dark purple varied with green, clothed with white hairs in front, rather broader than the chest: eyes dark red, rather large: feelers fourteen-jointed, black, clavate, hairy, much longer than the chest; first joint green, yellow towards the base; second cup-shaped; third and fourth very small; the following joints to the eleventh successively increasing in breadth and decreasing in length; club spindle-shaped, concave, broader than the eleventh joint and more than twice its length: chest spindle-shaped, coarsely shagreened, brassy green along the disk: fore-chest conical, abruptly narrower in front, much lower than the mid-chest; its length about half of its breadth: shield of the mid-chest nearly square, flat; sutures of the parapsides quite obsolete; axillæ parted by rather less than one-fourth of the breadth of the chest; scutcheon subrhomboidal, with a suture along each side: hind-chest and petiole very short: abdomen lance-shaped, brassy green, concave on the disk, slightly keeled beneath, hairy towards the tip, about twice the length and half the breadth of the chest; metapodeon occupying about one-sixth of the back; the following segments of nearly equal length: legs tawny; hips green; tips of feet brown: fore-legs short and stout with piceous thighs and shanks; middle-legs dilated as usual: wings tawny; veins also tawny; ulna much more than half the length of the humerus; radius a little shorter than the ulna; cubitus about half the length of the ulna, slightly curved; wing-brand very small. Length of the body 2 lines; of the wings 3 lines.

METALLON (n. g.)

Body narrow: head a little broader than the chest: feelers 10-jointed; first joint long, slender; second cup-shaped; third and following joints broader and successively decreasing in length: fore-chest of moderate size, short-conical, slightly wrinkled across the fore-part; its length about one-half of its breadth: shield of the mid-chest short; axillæ parted by a very short interval on the back; scutcheon subrhomboidal: hind-chest of moderate size, obconical, slightly declining: petiole very short: abdomen convex, smooth, elliptical; metapodeon short, hind-border slightly convex; octoon much longer; ennaton and decaton much shorter; paratelum and telum short.

METALLON *Acacallis*. Fem. *Fulvus, viridi varius, abdomine cupreo, antennis nigris apice fulvis, pedibus fulvis* Page 129.

Body tawny : eyes and eyelets dark red : head smooth, shining ; crown tinged with green : feelers black, clavate, hairy, tawny at the tips and towards the base, a little shorter than the chest ; the four basal joints, the fifth joint beneath, and the tenth joint are tawny : chest convex, smooth, shining, with a few punctures ; axillæ green ; scutcheon with a green line along the middle : abdomen shorter but hardly broader than the chest, copper-colour, tawny towards the base which is bright green : legs tawny ; hind-feet pale yellow with piceous tips : wings rudimentary. Length of the body $\frac{3}{4}$ line.

ENCYRTUS *Alycæus*. Fem. *Viridis, abdomine nigro-æneo, antennis femoribusque nigris, tibiis piceis. tarsis flavis, alis subgriseis.* Page 135.

Body short, broad : head and chest convex, dark green, punctured, rather hairy : head as broad as the chest : eyes dark red : feelers black, clavate, not longer than the chest ; first joint long, slender, slightly spindle-shaped : abdomen depressed above, smooth, shining, bronze-black, dark green at the base, much shorter but not narrower than the chest ; its length not exceeding its breadth : legs black ; shanks piceous ; knees and feet dull yellow ; tips of the latter piceous : wings slightly tinged with gray ; veins piceous. Length of the body $\frac{1}{3}$ line ; of the wings $\frac{2}{3}$ line.

ENCYRTUS *Molos*. Mas. *Cyaneo-viridis, abdomine æneo, antennis pedibusque nigris, tarsis flavis, alis limpidis.* Page 135.

Head and chest bluish green, convex, shagreened : head as broad as the chest : eyes dark red : feelers black, nearly filiform, rather broad, much longer than the chest ; first joint long, slender ; second cup-shaped ; the following joints nearly equal in length ; club spindle-shaped, full twice the length of the eighth joint : shield of the middle-chest long ; scutcheon also long, nearly obconical : abdomen dark bronze colour, smooth, shining, depressed above, narrower than the chest and hardly more than half its length : legs black ; knees and feet yellow ; tips of the latter piceous ; fore-feet tawny : wings colourless ; veins piceous. Length of the body $\frac{1}{2}$ — $\frac{3}{4}$ line ; of the wings 1 — $1\frac{1}{2}$ line.

ENCYRTUS Phrosime. Fem. *Ater, capite viridi, oviductu abdominis dimidii longitudine, tarsis flavis, protibiis et protar-sis fulvis, alis limpidis.* Page 135.

Head and chest convex: head green, as broad as the chest; crown narrow: eyes piceous, rather large: feelers black, clavate, not longer than the chest: chest broad, of a velvet-like black colour: abdomen black, much compressed, pointed, not more than one-half of the length of the breadth of the chest: sheaths of the oviduct projecting beyond the tip of the abdomen for about half its length: legs black; fore-shanks and fore-feet tawny; four hinder-feet pale yellow with brown tips: wings colourless; humerus tawny; cubitus yellow, forming a very acute angle with the radius, which like the ulna is almost black. Length of the body $\frac{1}{2}$ line; of the wings 1 line.

ENCYRTUS Tylissos. Mas. *Fuscus, pedibus fulvis, alis limpidis perangustis.* Page 135.

Body brown: head very short, as broad as the chest, which is rather flat: eyes piceous: feelers brown, hairy, filiform, nearly as long as the body; first joint stout, almost spindle-shaped: abdomen linear, flat, a little paler and longer but not narrower than the chest: legs tawny: wings colourless, very narrow; veins tawny. Length of the body $\frac{1}{3}$ line; of the wings $\frac{2}{3}$ line.

ENCYRTUS Iophon. Mas. *Niger, tarsis fulvis, alis limpidis.* Page 135.

Body black, stout: head a little narrower than the chest: eyes piceous: feelers black, subclavate, longer than the chest; first joint long, slender; second cup-shaped; the following joints from the fourth successively decreasing in length: chest broad, coarsely punctured: abdomen nearly linear, smooth, much shorter and narrower than the chest: legs black; knees and feet tawny: wings colourless; veins tawny. Length of the body $\frac{1}{2}$ line; of the wings 1 line.

ENCYRTUS Acratos. *Niger, abdomine nigro-æneo, pedibus piceis, alis limpidis.* Page 135.

Body black, short, broad: head nearly as broad as the chest which is slightly convex: eyes piceous, rather large: feelers black,

slender, subclavate, not longer than the chest ; first joint long, slender ; second cup-shaped ; the following joints successively decreasing in length ; club spindle-shaped, a little broader than the eighth joint and nearly thrice its length : head and chest dull, punctured : abdomen depressed, smooth, shining, bronze-black, much shorter but hardly narrower than the chest ; its length scarcely exceeding its breadth : legs piceous : wings colourless ; veins tawny. Length of the body $\frac{1}{3}$ line ; of the wings $\frac{2}{3}$ line.

ENCYRTUS Hydramon. Fem. *Viridis, abdomine aeneo, antennis pedibusque nigris, tarsis fulvis, alis limpidis.* Page 135.

Head and chest convex, shagreened, bright green : head a little broader than the chest : eyes and eyelets red : feelers black, clavate, hardly longer than the chest ; first joint long, slender, green ; second cup-shaped ; the following joints from the third to the eighth successively decreasing in length ; club spindle-shaped, a little broader than the eighth joint and more than thrice its length : chest short-elliptical : shield of the mid-chest long, conical ; axillæ just meeting on the back ; scutcheon large, subrhomboidal : abdomen obconical, depressed, shagreened, dark bronze colour, greenish towards the base, a little narrower and very much shorter than the chest : legs black ; feet dark tawny ; middle-feet and four hinder-knees pale yellow : wings colourless ; veins piceous. Length of the body $\frac{2}{3}$ line ; of the wings $1\frac{1}{3}$ line.

ENCYRTUS Philotis. Mas. *Niger, antennis tarsisque fulvis, alis limpidis.* Page 135.

Body black, rather narrow : head and chest convex, thickly punctured, rather hairy : head as broad as the chest : eyes and eyelets piceous : feelers dull tawny, filiform, rather hairy, nearly as long as the body ; first joint long, slender, black ; second cup-shaped, black ; the following joints from the third to the eighth long, linear, of nearly equal length ; club spindle-shaped, nearly twice the length of the eighth joint : chest nearly elliptical : shield of the mid-chest long ; axillæ parted by a short interval on the back ; scutcheon nearly obconical : abdomen obconical, depressed, smooth, a little narrower and very much shorter than the chest ; segments of nearly equal size : legs black ; knees black ; knees and feet dark tawny, middle-feet pale tawny : wings colourless ; veins piceous. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

ENCYRTUS Idmon. Fem. *Niger, tarsis fulvis, alis limpidis.*
Page 135.

Black: head and chest convex, shining, coarsely punctured, clothed with a few scattered hairs: head as broad as the chest: eyes dark red: feelers black, clavate, a little longer than the chest; first joint long, slender; second cup-shaped; the following joints from the third to the eighth short, successively and rapidly increasing in breadth; club elliptical, very much broader than the tenth joint and full thrice its length: chest short-elliptical: shield short and broad; axillæ nearly meeting on the back; scutcheon obconical; abdomen spindle-shaped, depressed, smooth, a little narrower and very much shorter than the chest: legs black; feet dark tawny, with piceous tips: wings colourless; veins piceous. Length of the body $1\frac{1}{2}$ line; of the wings 2 lines.

ENCYRTUS Alcomon. Fem. *Nigro-viridis, abdomine nigro-æneo, antennis pedibusque fulvis, alis subfulvis.* ENCYRTUS ænescens? Zetterstedt, *Ins. Lapp.*

Body narrow: head and chest convex, greenish black, with a bronze tinge, almost smooth: eyes and eyelets piceous: feelers sub-clavate, tawny, a little longer than the chest; first joint short and stout; second also short; third long; fourth and following joints short and successively decreasing in length; club conical: chest elliptical: shield rather flat; axillæ meeting on the back; scutcheon subrhomboidal: hind-chest short: abdomen elliptical, brassy black, flat, shining, a little shorter than the chest: legs piceous; middle-feet pale tawny; fore-legs and hind-feet dark tawny: wings slightly tawny, rather short and narrow; veins yellow; ulna and radius piceous. Length of the body $\frac{1}{2}$ line; of the wings $\frac{3}{4}$ line.

ENCYRTUS Genetyllis. Fem. *Nigro-æneus, capite abdominisque basi viridibus, antennis pedibusque nigris, tarsis fulvis, alis limpidis.* Page 135.

Body rather narrow: head and chest convex: head dark green, as broad as the chest: eyes dark red: feelers black, thick, hairy, clavate, shorter than the chest; first joint stout, spindle-shaped; second long cup-shaped; the following joints very short, but with a successive increase in breadth and length; club elliptical, very much broader than the preceding joint, and about five times its length: chest nearly linear, brassy-black, coarsely punctured; axillæ quite

meeting on the back : abdomen long-obconical, depressed, smooth, shining, dark bronze, bright green at the base, a little narrower but hardly shorter than the chest : legs black ; knees yellow ; feet tawny ; middle-shanks and middle-feet luteous : wings colourless, rather narrow ; veins tawny ; ulna piceous. Length of the body $1\frac{1}{2}$ line ; of the wings 1 line.

ENCYRTUS Anthores. Fem. *Nigro-viridis, antennis fulvis, pedibus nigris, tarsis luteis, alis fuscis.* Page 135.

Allied to *Epicopterus choreiformis*, *Eunotus cretaceus*, *Encyrtus encopiformis* and *E. (Ænasilus) Hyettus*. Body greenish black, very stout and short : head hardly broader than the chest, very coarsely excavated : eyes piceous : feelers tawny, slender, subelavate, a little longer than the chest ; first joint long, slender, dull yellow, traversed by a broad piceous band ; second piceous ; third small ; the following joints successively increasing in breadth ; club spindle-shaped, flat, a little broader than the preceding joint and about twice its length : chest broad, thickly punctured : fore-chest very short, but visible above : shield of the mid-chest very short ; its length less than one-half of its breadth ; scutcheon short-obconical ; axillæ parted from each other by a short interval : abdomen obconical, depressed, thinly punctured, a little shorter but not narrower than the chest ; its length slightly exceeding its breadth : legs deep black : feet luteous ; shanks slightly curved : fore-wings brown, hairy, broad, rather short ; veins piceous ; radius a little longer than the ulna ; cubitus a little longer than the radius, with which it forms a very acute angle : hind-wings colourless, narrow. Length of the body $\frac{1}{2}$ line ; of the wings $\frac{3}{4}$ line.

ENCYRTUS Idya. Fem. *Niger, capite nigro-æneo, antennis nigro-piceis fulvo cinctis, tibiis intermediis tarsisque fulvis, alis limpidis.* Page 135.

Body short : head and chest convex : head dark copper-colour, finely punctured, as broad as the chest : eyes and eyelets piceous ; the former rather large : feelers clavate, a little longer than the chest ; first joint black, long, spindle-shaped ; second long cup-shaped, piceous ; the following joints to the seventh tawny, very short ; club spindle-shaped, black, as long as the six preceding joints : chest short-oval, black, coarsely punctured : fore-chest short, visible above ; its length about one-fourth of its breadth : shield of the mid-chest short, broad ; axillæ nearly meeting on the back ; scutcheon nearly obconical : hind-chest extremely short : abdomen

oval, black, smooth, depressed above, deeply keeled beneath, a little shorter but hardly narrower than the chest : legs black ; feet and middle-shanks tawny : wings colourless ; veins tawny, piceous towards the tips. Length of the body $\frac{2}{3}$ line ; of the wings $1\frac{1}{4}$ line.

ENTEDON Isander. Mas. *Niger, antennis pedibusque flavis, alis limpidis.* Page 140.

Body black, slender, shining : head and chest convex ; head a little broader than the chest : eyes and eyelets piceous : feelers filiform, very slender, pale yellow, a little darker towards the tips, not much shorter than the body ; first joint long and slender ; second a little thicker ; third and following joints long and linear : chest elliptical : petiole short : abdomen depressed, a little narrower and shorter than the chest : legs pale yellow, long and slender ; hind-thighs piceous : wings colourless, rather long and narrow ; veins pale tawny ; ulna longer than the humerus ; radius and cubitus extremely short. Length of the body $\frac{1}{2}$ line ; of the wings $\frac{3}{4}$ line.

ENTEDON Nurscia. Mas. *Cyaneo-viridis purpureo varius, abdomine æneo basi viridi, antennis piceis, pedibus luteis, alis limpidis.* Page 140.

Head and chest convex, almost smooth : head green, short, impressed in front, a little broader than the chest : eyes and eyelets red : feelers piceous, filiform, a little longer than the chest ; first joint tawny, dilated, with a piceous line above ; second small ; third and following joints long, linear, of nearly equal size : chest bluish green with a purple tinge : sutures of the parapsides very distinct and deeply marked ; axillæ parted by one-third of the breadth of the chest ; scutcheon subrhomboidal, rather long : hind-chest of moderate size, obconical, declining, ridged : petiole not very long : abdomen nearly oval, dark bronze, smooth, depressed, bright green at the base, narrower and much shorter than the chest ; metapodeon occupying about one-third of the back ; octoon and following segments of moderate and nearly equal size : legs luteous ; hips green ; wings colourless or with a very slight brown tinge ; veins piceous ; ulna full twice the length of the humerus ; radius about one-third of the length of the ulna ; cubitus very short, not half the length of the radius. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

ENTEDON Sauros. Fem. *Viridis, micans, cyaneo æneoque varius, antennis nigris, pedibus flavis, alis limpidis.* Page 140.

Brilliant green: head and chest convex, finely shagreened, with a slight bluish tint: head very short, impressed in front, hardly so broad as the chest: eyes and eyelets red: feelers black, filiform, much shorter than the chest; first joint long, tawny towards the base; second cup-shaped; the following joints successively decreasing in length; club pointed, longer than the preceding joint: chest long-elliptical: fore-chest hardly seen above: shield of the mid-chest long; sutures of the parapsides distinct: axillæ parted by half the breadth of the chest; scutcheon subrhomboidal, with a slight suture on each side; these sutures converge together and form an angle near the tip: hind-chest rather long, obconical, declining, nearly smooth; its fore-shield distinct: petiole very short: abdomen elliptical, smooth, with a brassy tinge, depressed above, hardly keeled beneath, much shorter but hardly narrower than the chest; segments of nearly equal size: legs very pale yellow; hips green: wings colourless; fore-wings broad; veins piceous; ulna rather more than twice the length of the humerus; radius full half the length of the ulna, reaching very near the tip of the wing; cubitus very short; wing-brand small. Length of the body $\frac{2}{3}$ line; of the wings $1\frac{1}{3}$ line.

ENTEDON Axia. Mas. *Viridis, capite cyaneo, abdomine æneoviridi, antennis nigris, pedibus flavis, alis limpidis.* Page 140.

Head and chest convex, beset with a few scattered bristles: head thick, blue, almost smooth, a little broader than the chest, not impressed in front: eyes and eyelets red: feelers black, filiform, very much shorter than the chest; first joint long, slender, pale tawny, white at the tip; second long cup-shaped; third and following joints short; club ending in a bristle, about twice the length of the preceding joint: chest green, oval, well developed, very finely shagreened: fore-chest very short: shield of the mid-chest rather roughly shagreened; sutures of the parapsides very distinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal, rather large: hind-chest large, bluish green, obconical, declining, roughly excavated: petiole long, slender, finely punctured, a little longer than the hind-chest: abdomen æneous green, smooth, depressed, a little longer than broad; metapodeon occupying about one-fourth of the back; octoon and ennaton of moderate

size; decaton and protelum short; paratelum long; telum very small: legs bright yellow; hips green: wings colourless; forewings broad; veins tawny; ulna about twice the length of the humerus; radius not one-sixth of the length of the ulna; cubitus shorter than the radius: wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

ENTEDON Antaradus. Mas. *Niger, abdomine nigro-æneo basi cyaneo-viridi, antennis pedibusque nigris, tarsis fulvis, alis limpidis.* Page 140.

Body short, black: head and chest convex: head very short: eyes and eyelets red: feelers black, moniliform, very slender, longer than the chest; first joint very long and slender; second and following joints elliptical, slightly petiolated; club slender, pointed, much longer than the fifth joint: chest smooth, very short, hardly broader than long: fore-chest not visible above: petiole very short: abdomen bronze-black, depressed, smooth, nearly conical, bluish green at the base, a little longer and narrower than the chest: legs black, slender; feet tawny: wings colourless; forewings very broad; veins tawny; ulna much longer than the humerus; radius and cubitus extremely short; wing-brand very small. Length of the body $\frac{1}{3}$ line; of the wings $\frac{3}{4}$ line.

ENTEDON Erxias. Mas. *Niger, abdomine basi cyaneo-viridi, antennis pedibusque piceis, tarsis flavis, alis limpidis.* Page 140.

Body short: head and chest black, smooth, convex: head very short, impressed in front, as broad as the chest: eyes and eyelets dark red: feelers piceous, setaceous, a little longer than the chest; first joint long, stout, tawny: chest elliptical: petiole very short: abdomen short-elliptical, black, depressed, shorter but hardly narrower than the chest, bluish green at the base: legs piceous; feet and tips of the shanks yellow: wings colourless; veins pale tawny; ulna much longer than the humerus; radius and cubitus extremely short. Length of the body $\frac{1}{4}$ line; of the wings $\frac{1}{2}$ line.

ENTEDON Nænia. Mas. *Æneus, abdomine viridi-æneo, antennis nigris, pedibus flavis, alis subfulvis.* Page 140.

Body brass colour: head and chest convex, finely shagreened: head very short, impressed in front, hardly broader than the chest:

eyes and eyelets red: feelers black, slender, hairy, filiform, a little shorter than the chest; first joint long and slender, pale tawny towards the base; second cup-shaped; the following joints linear, rather long; club pointed, nearly twice the length of the last joint: chest oval: fore-chest extremely short: shield of the mid-chest less finely shagreened than are the other segments; sutures of the parapsides very distinct; axillæ parted by less than one-third of the breadth of the chest: scutcheon subrhomboidal, rather long: hind-chest obconical, declining, rather large, nearly smooth, with a ridge along the middle and a rim on each side: petiole very short: abdomen depressed, smooth, greenish brass colour, increasing in breadth from the base till near the tip, rather shorter and narrower than the chest; metapodeon occupying less than one-third of the back; octoon and following segments of nearly equal size: legs yellow; hips brassy; tips of the feet brown: wings slightly tawny; veins tawny; ulna rather more than twice the length of the humerus; radius not one-fourth of the length of the ulna; cubitus very short, not half the length of the radius: wing-brand extremely small. Length of the body 1 line; of the wings 2 lines.

ENTEDON Stennos. Mas. *Æneo-viridis, cupreo-purpureoque varius, antennis nigris, pedibus luteis, femoribus piceis, alis limpidis.* Page 140.

Head and chest convex, shagreened, bronze-green, varied with copper-colour and purple; the latter colours predominate towards the head and the fore-part of the chest: head thick, not impressed in front, a little broader and more finely shagreened than the chest; crown broad: eyes and eyelets dark red: feelers setaceous, black, slender, much shorter than the chest; first joint long, rather stout; second cup-shaped; the following joints slightly petiolated; club pointed, very much longer than the preceding joint: chest long-oval, well developed: fore-chest very short; its length not more than one-tenth of its breadth: sutures of the parapsides distinct, especially towards the hind border of the shield; axillæ parted by nearly one-half of the breadth of the chest; scutcheon subrhomboidal: hind-chest long-obconical, declining, with a ridge along the back and a rim on each side: petiole shagreened, stout, nearly one-third of the length of the abdomen, greenish brass colour, depressed, smooth, nearly conical, hardly more than half the length of the chest; metapodeon long; the following segments of moderate length: legs luteous; hips brass colour; thighs piceous; fore-shanks, fore-feet and tips of four hinder-feet tawny: wings colourless; veins piceous; ulna nearly twice the length of the humerus;

radius less than one-third of the length of the ulna; cubitus not half the length of the radius; wing-brand extremely small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

ENTEDON Amadocus. Fem. *Cyaneo-viridis, scutello purpureo-æneo, antennis nigris, tibiis tarsisque flavis, alis subfuscis.*
Page 140.

Head and chest convex, dark bluish green: head very finely shagreened, slightly impressed in front, hardly broader than the chest: eyes and eyelets dark red: feelers black, slightly setaceous, not longer than the chest, rather hairy; first joint long; second very small; the following joints successively decreasing in length; club conical, pointed, much longer than the preceding joint: chest long-oval: fore-chest not visible above: shield of the mid-chest rather long; sutures of the parapsides distinct towards the hind border of the shield; axillæ parted by nearly one-half of the breadth of the shield; scutcheon purplish bronze colour, nearly obconical: hind-chest well developed, obconical, declining; its fore-shield as long as the following segment: petiole very short: abdomen bluish green, smooth, rather hairy, depressed above, hardly keeled beneath, much shorter and a little narrower than the chest; metapodeon occupying nearly two-thirds of the back: octoon and ennaton of moderate size; the following segments very short: legs yellow; hips and thighs bluish green; fore-feet and tips of four hinder-feet tawny: wings slightly brown; veins piceous; ulna rather more than twice the length of the humerus; radius full one-third of the length of the ulna; cubitus much less than half the length of the radius, but longer than is usual in this genus; wing-brand of moderate size. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{3}{4}$ line.

ENTEDON Temena. Mas. *Viridis, capite aureo-viridi, abdomine cyaneo-æneo basi viridi, antennis nigris, pedibus albis, alis limpidis.* Page 140.

Head and chest convex: head bright golden green, smooth, not impressed in front, bluish green behind, a very little broader than the chest: eyes and eyelets red: feelers black, filiform, not more than half the length of the chest; first joint long, rather slender; second long cup-shaped; the following joints linear, rather long; club pointed, much longer than the preceding joint: chest green, long-oval, well developed: fore-chest very short, but distinct: sutures of the parapsides deeply marked; axillæ parted by rather more than one-third of the breadth of the chest; scutcheon sub-

rhomboidal, rather long and large : hind-chest long, obconical, declining, with a ridge down the middle and a rim on each side ; fore-shield very short : petiole slender, cylindrical, shagreened, almost as long as the hind-chest : abdomen bluish bronze, depressed, smooth, a little longer than broad, narrower than the chest and hardly half its length ; metapodeon green, occupying full one-third of the back ; octoon and following segments very short : legs white ; hips green ; tips of the feet brown : wings colourless ; veins dark tawny ; ulna about twice the length of the humerus ; radius one-fourth of the length of the ulna ; cubitus not half the length of the radius ; wing-brand very small. Length of the body $\frac{3}{4}$ line ; of the wings $1\frac{1}{2}$ line.

ENTEDON Metagenes. Mas. *Purpureo-cyaneus, capite scutellorque viribibus, abdomine nigro basi viridi, antennis pedibusque nigris, tarsis fulvis, alis limpidis.* Page 140.

Body smooth, shining : head and chest convex : head green, very short, impressed in front, as broad as the chest : eyes and eyelets dark red : feelers black, nearly filiform, hardly longer than the chest ; first joint long, rather slender, slightly curved ; second somewhat dilated ; the following joints long, linear : club pointed at the tip, much longer than the preceding joint : chest elliptical, very dark blue, tinged with purple ; scutcheon green : petiole very short : abdomen short-elliptical, depressed, black, green at the base, a little shorter and broader than the chest : legs black ; feet tawny ; tips piceous : wings colourless ; veins piceous ; ulna nearly twice the length of the humerus ; radius and cubitus extremely short. Length of the body $\frac{1}{2}$ line ; of the wings 1 line.

ENTEDON Ergetelis. Fem. *Cyaneo-viridis, capite aureo-viridi, abdomine æneo, antennis nigris, pedibus albis, alis limpidis.* Page 140.

Head and chest convex : head golden green, very finely shagreened, slightly impressed in front, hardly as broad as the chest : eyes red, rather large : feelers black, filiform, much shorter than the chest ; first joint pale tawny ; second rather short ; the following joints long, but successively decreasing in length ; club pointed, longer than the preceding joint : chest elliptical, bluish green, rather coarsely shagreened : fore-chest very short, just visible above : sutures of the parapsides rather distinct ; axillæ parted by half the breadth of the chest ; scutcheon subrhomboidal ; sutures inclined towards each other near the tip, but not meeting : hind-chest short :

petiole very short: abdomen smooth, depressed, bronze colour, hardly keeled beneath, shorter but scarcely narrower than the chest: segments of nearly equal size: legs white; hips green: wings colourless; veins piceous; ulna full twice the length of the humerus; radius about one-third of the length of the ulna; cubitus one-fourth of the length of the radius; wing-brand small. Length of the body $\frac{1}{2}$ line: of the wings 1 line.

ENTEDON Adreus. Mas. *Cyaneus, abdomine viridi, antennis nigris, pedibus flavis, femoribus viridibus, alis limpidis.*
Page 140.

Head and chest convex, blue, finely shagreened: head broader than the chest, short, impressed in front: eyes and eyelets dark red: feelers black, nearly moniliform, hardly longer than the chest; first joint long, dilated; second short; third and following joints longer; club long, pointed, much longer than the preceding joint: chest elliptical: sutures of the parapsides distinct: petiole rather long: abdomen dark green, convex, smooth, shorter and narrower than the chest: legs pale yellow; hips and thighs green; tips of feet brown; fore-shanks and fore-feet tawny: wings colourless; veins tawny; ulna nearly twice the length of the humerus; radius hardly one-half the length of the ulna; cubitus short, but longer than usual in the genus. Length of the body $\frac{1}{2}$ line: of the wings 1 line.

ENTEDON Ictas. Mas. *Æneo-viridis, abdomine æneo, antennis nigris, pedibus flavis, femoribus fulvis, alis subfulvis.*
Page 141.

Head and chest convex: head bright green, very short, impressed in front, very finely shagreened, a little broader than the chest: eyes and eyelets red: feelers black, setaceous, shorter than the chest; first joint long, rather stout; second long cup-shaped; the following joints slightly petiolated; club pointed, ending in a bristle, much longer than the preceding joint: chest elliptical, brassy green, finely shagreened: fore-chest extremely short: shield of the mid-chest short, roughly shagreened; sutures of the parapsides very distinct; axillæ parted by one-third of the breadth of the chest; scutcheon subrhomboidal, long: hind-chest long, obconical, declining: petiole very short: abdomen copper-colour, smooth, depressed, slightly increasing in breadth from the base till near the tip, shorter but hardly narrower than the chest; metapodeon rather long; octoon and following segments of moderate size: legs pale

yellow; hips brassy; thighs tawny: wings slightly tawny; veins tawny; ulna about twice the length of the humerus; radius not near one-fourth of the length of the ulna; cubitus very short; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

EULOPHUS Mamurius. Mas. *Viridis, abdomine nigro-æneo basi viridi, antennis nigris, tarsis piceis basi flavis, alis subgriseis.* Page 144.

Head and chest green: head a little broader than the chest: eyes and eyelets red: feelers black, hairy, 3-branched, subclavate, a little longer than the chest; first joint green, spindle-shaped: chest long-elliptical, very finely shagreened: shield of the mid-chest rather flat; sutures of the parapsides indistinct; axillæ parted by full one-third of the breadth of the chest: hind-chest nearly smooth, obconical, declining; its fore-shield of moderate length: abdomen linear, depressed, bronze-black, green at the base, narrower but not shorter than the chest: legs green; feet piceous; knees and base of four hinder-feet yellow: wings slightly gray; veins tawny; ulna a little shorter than the humerus; radius less than one-half of the length of the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings 1 line.

EULOPHUS Rhamnius. Mas. *Viridis, capite cyaneo, scutello nigro, abdomine nigro-æneo, basi fulvo maculato, antennis pedibusque nigris, tibiis anticis tarsisque piceis, alis subgriseis.* Page 144.

Head blue, as broad as the chest: eyes and eyelets piceous: feelers black, hairy, subclavate, a little longer than the chest; first joint spindle-shaped: chest long-elliptical, dark green, shagreened: fore-chest very short: shield of the mid-chest broad; sutures of the parapsides indistinct; axillæ parted by one-third of the breadth of the chest; scutcheon subrhomboidal, black: hind-chest rather large, obconical, declining, finely shagreened, with a ridge along the back: petiole very short: abdomen shorter and much narrower than the chest, very slightly increasing in breadth from the base to near the tip, depressed, smooth, bronze-black, with a dark tawny spot near the base which is green: legs black; knees, feet, and fore-shanks piceous: wings slightly gray; ulna full as long as the humerus; radius more than half the length of the ulna; cubitus less than one-half the length of the radius; wing-brand small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

EULOPHUS stygius. Fem. *Niger, pedibus fulvis, femoribus nigris, alis fuscis*. Page 144.

Black, hairy, very finely shagreened, almost smooth: head convex: eyes and eyelets red: feelers black, clavate, hairy, shorter than the chest; first joint long, slender; second cup-shaped; the following joints successively decreasing in length; club conical, a little longer than the preceding joint; abdomen long-elliptical, smooth, depressed, hairy, much shorter and a little broader than the chest; metapodeon occupying nearly one-fourth of the back; the following segments of moderate size: legs tawny; hips and thighs black; tips of the latter tawny: wings brown; veins piceous; ulna much longer than the humerus; radius more than half the length of the ulna; cubitus not half the length of the radius; wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{2}$ lines.

EULOPHUS Pythodorus. Fem. *Æneus, capite viridi, abdomine purpureo basi viridi, antennis femoribusque nigris, tibiis piceis, tarsis flavis, alis limpidis*. Page 144.

Body short and broad: head and chest convex, very finely shagreened: head short, dark green, much impressed in front: eyes and eyelets dark red: feelers black, clavate, shorter than the chest; first joint long, slender; second cup-shaped; three following joints successively decreasing in length; club long, pointed, full twice the length of the preceding joint: chest bronze colour, short-elliptical: fore-chest very short: shield of the mid-chest broad; sutures of the parapsides indistinct; axillæ parted by nearly half the breadth of the chest; scutcheon subrhomboidal: hind-chest short, transverse, declining, with a ridge along the middle: petiole very short: abdomen short-elliptical, dark purple, very finely shagreened, green at the tip, bright green at the base, depressed above, hardly keeled beneath, a little shorter and broader than the chest; metapodeon occupying full one-sixth of the back; the following segments of moderate and nearly equal size: hips and thighs black; shanks piceous; feet yellow with piceous tips; fore-shanks and fore-feet tawny: wings colourless; humerus longer than the ulna; radius less than half the length of the ulna; cubitus shorter than the radius; wing-brand small. Length of the body 1 line; of the wings 2 lines.

EULOPHUS Acalles. Mas. *Cyaneus, abdomine purpureo-æneo flavo-maculato basi viridi, antennis nigris, pedibus albis, femoribus nigris, tibiis apice piceis, alis limpidis.* Page 144.

Head and chest blue: feelers black, hairy, a little shorter than the chest: shield of the mid-chest flat; scutcheon black: hind-chest nearly smooth, obconical, declining; its fore-shield of moderate length: abdomen nearly linear, depressed, smooth, purplish bronze, yellow towards the base which is green: legs white; hips and thighs black; tips of the shanks piceous; fore-shanks and fore-feet tawny: tips of the four hinder-feet brown: wings colourless; fore-wings broad; veins tawny; ulna shorter than the humerus; radius much shorter than the ulna; cubitus about one-third of the length of the radius; wing-brand small. Length of the body 1 line; of the wings $1\frac{3}{4}$ line.

EULOPHUS Veturius. Mas. *Viridis, abdomine purpureo-æneo basi viridi, antennis tibiisque piceis, tarsis fulvis basi flavis, alis limpidis.* Page 144.

Green: head broader than the chest: eyes and eyelets dark red: feelers piceous, 3-branched, a little longer than the chest; first joint green, spindle-shaped: chest oval, finely shagreened: shield of the mid-chest broad, rather short; sutures of the parapsides indistinct; scutcheon subrhomboidal: hind-chest nearly smooth, obconical, declining, its fore-shield of moderate length: abdomen linear, depressed, purplish bronze, smooth, green at the base, hardly shorter but very much narrower than the chest: hips and thighs green; shanks piceous; knees and feet tawny; base of four hinder-feet pale yellow: wings colourless; fore-wings rather broad; veins tawny; ulna a little longer than the humerus; radius less than half the length of the ulna; cubitus much shorter than the radius; wing-brand very small. Length of the body $\frac{3}{4}$ line; of the wings $1\frac{1}{2}$ line.

EULOPHUS Scyllis. Fem. *Viridis, abdomine æneo, antennis nigris, pedibus fulvis, alis subfulvis.* Page 144.

Head and chest green, convex, finely shagreened: head as broad as the chest: eyes and eyelets red: feelers black, clavate, shorter than the chest; first joint long, slender; second cup-shaped; the four following joints successively decreasing in length; club

conical, longer than the preceding joint: chest long-elliptical: fore-chest rather long, short-conical, its length full one-half of its breadth: sutures of the parapsides very distinct; axillæ parted by one-fourth of the breadth of the chest; scutcheon subrhomboidal: hind-chest well developed, obconical, declining, almost smooth, with a ridge along the back; its fore-shield rather long: petiole very short: abdomen bronze colour, slightly tinged with green, smooth, elliptical, depressed above, slightly keeled beneath, about as long and as broad as the chest; metapodeon occupying about one-sixth of the back; the following segments of moderate size: legs tawny; hips green: wings slightly tawny; veins tawny; ulna as long as the humerus; radius much shorter than the ulna; cubitus not half the length of the radius; wing-brand very small. Length of the body $1\frac{1}{4}$ line; of the wings $2\frac{1}{4}$ lines.

DIGLYPHUS Poppœa. Fem. *Viridis, cyaneo purpureo æneoque varius, antennis nigris basi albis, pedibus viridibus albo-cinctis, alis limpidis.* Page 145.

Head blue, nearly smooth, bronze on the crown, impressed above, prominent in front, not broader than the chest: eyes and eyelets red: feelers black, stout, clavate, shorter than the chest; first joint long, slender, white towards the base; second of moderate size; third and fourth broader; club long-conical, twice the length of the preceding joint: chest green, elliptical: fore-chest extremely short: shield broad, nearly flat above; sutures of the parapsides obsolete; axillæ parted by nearly one-half of the breadth of the chest; scutcheon nearly obconical, with a suture along each side: hind-chest well developed, obconical, slightly declining, nearly smooth, with a slight ridge along the back; fore-shield large, about one-half of the length of the following segment: petiole very short: abdomen smooth, elliptical, purplish bronze, green at the base, blue at the tip, depressed above, slightly keeled beneath, shorter but hardly broader than the chest; metapodeon occupying full one-half of the back; oetoon and enuaton rather large; the following segments extremely short: hips and thighs green; shanks piceous with a green tinge; feet and tips of thighs and of shanks white; fore-shanks, fore-feet, and tips of four hinder-feet tawny: wings colourless; veins piceous; ulna nearly as long as the humerus; radius much shorter than the ulna; cubitus not half the length of the radius; wing-brand small. Length of the body $\frac{1}{2}$ line; of the wings $1\frac{1}{4}$ line.

DIGLYPHUS *Aculeo*. Fem. *Viridis, abdomine fulvo, antennis pedibusque flavo-fulvis, alis subfulvis.* Page 145.

Body very slender: head and chest green, rather flat: head broader than the chest: eyes and eyelets red: feelers pale tawny, clavate, a little shorter than the chest; club much broader than the preceding joint and more than twice its length: chest long-elliptical: petiole very short: abdomen elliptical, tawny, smooth, depressed, a little shorter but not narrower than the chest: legs pale tawny: wings narrow, nearly colourless or slightly tawny; veins tawny; ulna a little longer than the humerus; radius and cubitus very short. Length of the body $\frac{1}{2}$ line; of the wings $\frac{2}{3}$ line.

Var. β . Feelers and legs yellow.

TETRASTICHUS *Triarius*. Mas. *Niger, abdomine nigro-æneo, antennis fulvis, pedibus piceis, tarsis flavis, alis limpidis.* Page 151.

Body rather short: head and chest black, convex, smooth: head short, as broad as the chest: eyes and eyelets bright red: feelers tawny, adorned with long hairs, not longer than the chest; first joint piceous: chest elliptical: fore-chest very short: shield of the mid-chest broad and long, with a suture along the back; sutures of the parapsides distinct; axillæ parted by more than one-third of the breadth of the chest; scutcheon subrhomboidal: hind-chest transverse, declining: petiole very short: abdomen bronze-black, depressed, linear, shorter and narrower than the chest: legs piceous; four hinder-knees and feet pale yellow; tips of the latter brown; fore-shanks and fore-feet tawny: wings colourless; veins tawny; ulna longer than the humerus; radius next to none; cubitus about one-fourth of the length of the ulna; wing-brand extremely small. Length of the body $\frac{1}{3}$ line; of the wings $\frac{3}{4}$ line.

TETRASTICHUS *Abydenus*. Mas. *Niger, abdomine fulvo maculato, antennis fulvis, pedibus flavis, alis limpidis.* Page 151.

Head and chest black, convex, shining: head smooth, very short, hardly as broad as the chest: eyes and eyelets red; feelers filiform, tawny, slender, a little longer than the chest, adorned with long hairs; first joint piceous; second yellow: chest nearly oval, very finely punctured: fore-chest extremely short: shield of the mid-chest very long, with a slight suture along the middle; sutures of the parapsides strongly marked; scutcheon nearly obconical,

with a suture along each side : hind-chest rather large, obconical, declining, with a ridge along the middle ; its fore-shield tolerably long : petiole very short : abdomen black, smooth, shining, depressed, nearly linear, with a large tawny spot near the base, a little longer and much narrower than the chest : legs pale yellow ; hips black ; tips of feet tawny : wings colourless, ample ; veins tawny ; ulna a little longer than the humerus ; radius next to none ; cubitus not one-fourth of the length of the ulna ; wing-brand very small. Length of the body 1 line ; of the wings $2\frac{1}{4}$ lines.

TETRASTICHUS Bermius. Mas. *Niger, abdomine æneo, antennis fulvis, pedibus flavis, alis limpidis.* Page 151.

Head and chest black, convex, smooth, shining : head as broad as the chest : eyes and eyelets bright red : feelers slender, filiform, pale tawny, adorned with long hairs, nearly as long as the body ; first and second joints pale yellow : chest short : shield of the mid-chest broad ; sutures of the parapsides very distinct ; scutcheon short, with a suture along each side : abdomen bronze-colour, linear, depressed, shorter and narrower than the chest : legs pale yellow : wings colourless, deeply fringed ; veins tawny ; ulna nearly as long as the humerus ; radius next to none ; cubitus as usual ; wing-brand very small. Length of the body $\frac{1}{2}$ line ; of the wings $1\frac{1}{4}$ line.